

93.05.07

GREEN CROSS(R)

BUG KILLER

DOMESTIC

INSECTICIDE DUST

Controls insects on potatoes, cucumbers, beans, and other vegetables.

NET WEIGHT: 2 kg, 4 kg

GUARANTEE:

Carbaryl.....5%

READ THE LABEL BEFORE USING

REGISTRATION NO.: 7,652
PEST CONTROL PRODUCTS ACT

Fisons Horticulture Inc.
600 - 25 Watline Avenue
Mississauga, Ontario
L4Z 2Z1

DIRECTIONS:

Apply insecticide dust using a duster or shaker device.

Potatoes: To control Colorado potato beetles, flea beetles and leafhoppers, apply a light even coating on upper and lower leaf surfaces. For larger areas, apply 225 grams/100 square meters to control Colorado potato beetle and 450 grams/100 square meters to control flea beetles and leafhoppers. If necessary, repeat treatment at 7 - 10 day intervals.

Cucumbers And Squash: To control cucumber beetles, squash bugs

and flea beetles, dust upper and lower leaf surfaces evenly. For larger areas use 275 - 450 grams/100 square meters. If necessary, repeat at 7 - 10 day intervals.

Beans: To control Mexican bean beetle, leafhoppers and flea beetles, dust upper and lower leaf surfaces evenly. For larger areas use 225 - 450 grams/100 square meters. Repeat in 10 - 14 days as needed.

Sweet Corn:

Corn Borer: Make first application when egg masses on leaves are in black-head stage. Apply evenly on upper and lower leaf surfaces. A minimum of 4 applications are required at 4 - 5 day intervals. For larger areas, apply 450 - 675 grams/100 square meters.

Corn Earworm: Dust leaves and ears thoroughly and evenly at 2 - 3 day intervals starting at first silking and continuing until silks start drying. For larger areas apply 450-675 grams /100 square meters. Three or more applications are necessary depending upon the length of silking and the severity of the infestation.

*There is no danger of residue in cattle which eat treated corn.

TOMATOES: To control flea beetles, fall armyworm, stink bugs, tomato fruitworm and tomato hornworm, apply a light even coating to upper and lower leaf surfaces. For larger areas use 225-325 grams/100 square meters. Repeat at 7 - 10 day intervals as needed.

PRECAUTIONS:

KEEP OUT OF REACH OF CHILDREN. This product is toxic to bees. AVOID USING DURING FLOWERING PERIOD OF THE CROP. Do not breathe dust or take internally. Avoid skin contact. Change clothing after use. Avoid contamination of feed, feed troughs and water receptacles. Do not treat within 7 days of harvest except tomatoes and corn which may be treated 1 day before harvest.

Do not use on Boston ivy or Virginia creeper.

FIRST AID:

If in eyes or on skin, wash with plenty of water. If swallowed,

induce vomiting and obtain medical attention. Do not induce vomiting if patient is unconscious or in a convulsive state.

TOXICOLOGICAL INFORMATION:

Carbaryl is a moderate reversible cholinesterase inhibitor. Atropine is antidote.

DISPOSAL:

Dispose of empty container with household garbage.

Product Use Information: 1-800-668-5669

(R) Registered trademark of Fisons Horticulture Inc.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))