

31-JAN-2000

(leaflet)

HAGEN

FLEA COLLAR WITH INTEGRAL BUCKLE FOR MEDIUM DOGS

DOMESTIC

READ THE LABEL AND LEAFLET BEFORE USING.

GUARANTEE.....PROPOXUR 9.4%

REG. NO. 23541 P.C.P. ACT

NET CONTENTS: 30.2 g

DISTRIBUTED BY:
ROLF C. HAGEN INC.
3225 SARTELON
MONTREAL, QUEBEC H4R 1E8

PRINTED IN CANADA/MADE IN GERMANY

DIRECTIONS: For best results, before placing this collar on, bathe and shampoo pet with a pyrethrin based product such as Hagen Flea & Tick Shampoo for Dogs. This will produce a "quick kill" effect.

Remove the collar from the package and place around dog's neck. Thread flat end of the collar through buckle hole and tighten as necessary, so that at least two ridges are covered by the buckle, leaving ample room for expansion as the dog grows. Cut off any excess length and discard in garbage. Collar must be worn loosely, but securely enough to prevent easy removal. After collar placement, adult ticks will be killed in a few days and will fall off or may be easily removed. American Dog and Brown Dog ticks appear on dogs in three stages. In the first two stages, ticks are small and are difficult to see. The collar will also control these immature stages. Collar will kill ticks encountered in first 2 months of wear. Kills dog fleas for up to

4 months. FOR CONTINUOUS PROTECTION, REPLACE COLLAR WHEN EFFECTIVENESS DIMINISHES. To release the collar, squeeze between thumb and finger and pull sharply.

FLEAS: This collar is also effective in killing fleas. Temporary wetting will not reduce effectiveness. If your dog goes swimming or is out in the rain, it is not necessary to remove the collar. The continuous action collar will rapidly replace any flea and tick killing material removed. For best results, use in combination with a complete flea eradication program (consult your local pet retailer for more information).

PRECAUTIONS: Do not open protective pouch until ready to use. DO NOT ALLOW CHILDREN TO HANDLE THIS COLLAR. Dust will form on the collar during storage. Do not get dust or collar in the mouth. Harmful if swallowed. Do not get dust in eyes. Wash hands thoroughly with soap and water after handling. Do not use on dogs younger than three months old. When collar is first worn, observe neck area every few days for irritation or adverse reaction. Collar is intended for use only as an insecticide generator and is not to be taken internally by humans or animals. Do not use on sick or convalescing dogs, or dogs under drug treatment. Do not use other cholinesterase inhibitors (other organophosphate or carbamate containing parasitocides) while collar is worn or for a few days after removing collar. Before using on pregnant animals, consult a veterinarian.

FIRST AID: If dust gets in eyes or on skin, flush with plenty of water. If irritation develops or persists, or if ingested, obtain medical attention.

TOXICOLOGICAL INFORMATION: Propoxur is a cholinesterase inhibitor. Atropine is antidotal.

DISPOSAL: Wrap and dispose of any excess collar, packaging and expired collars with garbage to keep out of reach of children.

(box label)

HAGEN

CONTENTS: 1 COLLAR/NET WT. 30.2 g

FLEA COLLAR WITH INTEGRAL BUCKLE

FOR MEDIUM DOGS FITS DOGS WITH NECKS UP TO 52.5 cm.

3 STEP FLEA ATTACK PROGRAM
PREVENTATIVE TREATMENT

KILLS FLEAS FOR UP TO 4 MONTHS AND AMERICAN
DOG AND BROWN DOG TICKS UP TO 2 MONTHS.

DOMESTIC - READ THE LABEL AND ENCLOSED LEAFLET BEFORE USING.

GUARANTEE: 9.4% PROPOXUR

REG.NO. 23541 P.C.P. ACT

DISTRIBUTED BY:
ROLF C. HAGEN INC.
3225 SARTELON
MONTREAL, QUEBEC H4R 1E8

PRINTED IN CANADA/MADE IN GERMANY

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))