

King Logo

RODENTEX® Mini-Block Mini-Pack
With Bitrex®

Kills Rats and Mice with a single feeding
Kills Warfarin-Resistant Norway Rats
DOMESTIC

GUARANTEE

Bromadiolone0.005 %

REG. NO. 26915 PCP ACT

KEEP OUT OF REACH OF CHILDREN
READ THE LABEL BEFORE USING
WARNING, CONTAINS THE ALLERGENS PEANUTS AND WHEAT

WARNING



POISON

Net Contents : 84 g

King Home & Garden Inc.
3-304 Stone Road West,
Guelph, Ontario N1G 4W4
1-866-458-5464

DIRECTIONS FOR USE:

Use Rodentex® Mini-Blocks for the control of Norway rats, Roof rats and House mice indoors and against the outside walls of buildings.

USE LIMITATIONS: Bait must either be placed in tamper-resistant bait stations or in locations not accessible to children, pets or livestock. **DO NOT** place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food.

Selection of Treatment Areas: Place bait in locations where rats and mice travel and will most likely find and consume the bait. Placement along walls, in corners, by gnawed openings, in and around burrows, and in concealed locations where rats and mice feed is desirable. Maintain a steady supply of fresh bait. Remove any spoiled or contaminated bait promptly.

Application Directions:

For Norway Rats (*Rattus norvegicus*) and Roof Rats (*Rattus rattus*): For both species, eliminate as much of the naturally occurring food as possible and place 3 to 10 pieces of block (usually at intervals of 5-10 m) within the treatment area. Maintain an uninterrupted supply of fresh bait for at least 10 days, or until signs of rat activity cease.

For House Mice (*Mus musculus*): Eliminate as much of the naturally occurring food as possible and place 1 piece of block at intervals of 2-4 m within the treatment area. If a bait placement is not disturbed after several days, move the placement approximately 1 metre. Two (2) pieces of block per placement may be necessary at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 days or until signs of mouse activity cease.

Note: Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and unconsumed bait properly. To prevent re-infestation, eliminate food, water and harbourage as much as possible. If re-infestation occurs, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations as necessary.

PRECAUTIONS:

KEEP OUT OF REACH OF CHILDREN, PETS AND LIVESTOCK. May be harmful or fatal if swallowed or absorbed through the skin.

Rubber gloves must be worn when handling product and when disposing of dead rodents, unconsumed bait and empty containers. Avoid contact with eyes, skin and clothing. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Wash skin thoroughly with soap and water after handling. Wash contaminated clothing, separately from other laundry, with soap and hot water before reuse. **KEEP AWAY FROM FEED AND FOODSTUFFS.** Harmful or fatal, if swallowed or absorbed through the skin. This material may reduce the clotting ability of the blood and cause bleeding. This product is toxic to fish and wildlife. Keep out of lakes, streams or ponds.

FIRST AID:

If in eyes, flush with water for 15 minutes. If irritation persists, seek medical attention. **If on skin**, wash affected area thoroughly with soap and water. **If swallowed**, give patient one or two glasses of water and INDUCE VOMITING by touching back of throat with a finger or blunt object; repeat until vomitus is clear. DO NOT INDUCE VOMITING or give anything by mouth to an unconscious person. For all cases of human ingestion IMMEDIATELY notify a physician or poison control centre. Take container, label or product name and Pest Control Product Registration Number with you. If pet or livestock poisoning is suspected, immediately contact a veterinarian.

TOXICOLOGICAL INFORMATION:

Vitamin K₁ in the form of intramuscular or subcutaneous injections, or by oral ingestion are suggested remedial treatments for anticoagulant poisoning. The severity of the case measured by establishing prolonged prothrombin times (P.T.) will determine appropriate therapy. Monitoring P.T. will indicate the necessity of repeated treatments.

STORAGE AND DISPOSAL:

Store in a cool dry place away from other chemicals and food or feed. Store products not in use, in original container, in a secure location inaccessible to children and non-target animals. Do not contaminate water, food or feed by storage or disposal. Do not reuse empty container. Wrap empty container and dispose in household garbage.

RODENT DISPOSAL:

Dispose of dead rodents in garbage or by burying.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.