

89.12.21

(( CARTON ))

SENDRAN (R)

INTEGRAL BUCKLE

Flea & Tick Collar for Dogs

Kills Fleas & Ticks up to 4 Months

For continuous protection replace collar  
when effectiveness diminishes

SOLD THROUGH  
VETERINARIANS

CAUTION:  
DO NOT ALLOW CHILDREN TO PLAY WITH THIS COLLAR

DOMESTIC

READ LABEL BEFORE USING

GUARANTEE: Propoxur.....9.4%

CONTENTS: 1 Collar, Net Weight 32.7 g

Registration No. 18507 P.C.P. Act

Distributed by  
BAYVET Division  
Chemagro Limited  
Etobicoke, Ontario,  
Canada M9W 1G6

HAYER (R)

(( INSERT ))

SENDRAN (R)

INTEGRAL BUCKLE

Flea and Tick Collar  
For Dogs

DOMESTIC

GUARANTEE: Propoxur\* 9.4%

DIRECTIONS: Buckle collar around dog's neck. Allow a spacing of two fingers between collar and neck to prevent irritation and cut off any excess length. When collar is first placed on dog, adult ticks will be killed over entire body in a few days and will fall off or may then be easily removed. Ticks appear on dogs in three stages. In the first two stages, ticks are smaller than a match head and are difficult to see. The collar will also control these immature stages. If ticks are a problem, this collar should be worn constantly. KILLS FLEAS AND TICKS FOR UP TO 4 MONTHS. FOR CONTINUOUS PROTECTION, REPLACE COLLAR WHEN EFFECTIVENESS DIMINISHES.

Fleas - This collar is also highly effective in killing fleas. As long as your dog is wearing a SENDRAN Flea & Tick Collar, do not use a separate flea collar. Temporary wetting will not reduce effectiveness. If your dog goes swimming or is out in the rain, it is not necessary to remove the collar. The continuous action collar will rapidly replace and tick and flea killing material removed. Remove collar while dog is being bathed.

PRECAUTION: Do not open protective pouch until ready to use. Do not allow children to handle this collar. Dust will form on this collar during storage. Do not get dust or collar in mouth, harmful if swallowed. Do not get dust in eyes. Wash hands thoroughly with soap and water after handling. Do not use on dogs younger than 3 months old. Consult a veterinarian before use on pregnant animals. When collar is first worn observe neck area every few days for irritation. Remove collar at first sign of irritation or adverse reaction. Collar is intended for use only as an insecticide generator and is not to be taken internally by man or animals. Do not use on sick or convalescing dogs. Do not use other pesticides on dog or administer other cholinesterase inhibitors (e.g. oral parasiticides) while collar is worn or for a few days after removing collar. Dispose of any excess collar, packaging and expired collars with garbage to keep out of reach of children.

FIRST AID: In case of poisoning, call a physician immediately. If dust from collar contacts skin, wash thoroughly with soap and water. If dust contacts eyes, flush eyes with water and get medical attention.

TOXICOLOGICAL INFORMATION: Dust in eyes will cause temporary pupillary constriction\*. Dust released from collar is a cholinesterase inhibitor.

NOTE TO PHYSICIAN: Atropine and homatropine\* are antidotal.

SENDRAN (R) is a Reg. TM of the parent company of Farbenfabriken Bayer, GmbH, Leverkusen.

BAYVET Division CHEMAGRO Ltd. Rexdale, Ontario M9W 1G6

PRINTED IN CANADA

\*\*\*\*\*

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))