

85.12.31

----- CARTON -----

HAGEN (R)

4 plus 2

Flea and Tick Collar for Cats

4 plus 2

Flea and Tick Collar for Cats

Kills fleas for up to 4 months

Kills ticks for up to 2 months

DOMESTIC

Read label before using

GUARANTEE Propoxur 9.4%

Reg. No. 17743 P.C.P. Act.

HAGEN (R)

Art #C-142

Net Weight 8.35 g

Distributed by:  
Rolf C. Hagen Inc.,  
Montreal, Quebec  
H4R 1E8

----- INSERT -----

HAGEN (R)

HAGEN FLEA & TICK COLLAR FOR CATS

DIRECTIONS: Remove collar from package and place around cat's neck. Adjust for proper fit and buckle in place. Collar must be

worn loosely but securely enough to prevent easy removal and loss. Cut off any excess length of collar, leaving ample length for expansion as the cat grows and discard of by discarding in trash. Check and adjust collar periodically to assure a proper fit. Do not use on kittens under 12 weeks of age.

RESULTS EXPECTED: The fleas and tick collar starts killing fleas and ticks when it is placed around the cat's neck. Fleas will be killed and new ones which may temporarily appear on the cat, will also be killed during the 4 months. Non-engorged ticks are tough occasional pests of cats and will be killed within a few days. Collar will kill ticks encountered in first two months of wear. FOR CONTINUOUS PROTECTION REPLACE COLLAR WHEN EFFECTIVENESS DIMINISHES.

Some cats may be sensitive to this collar. When collar is first worn observe neck area every few days for irritation. Any collar, when fastened too tightly, may cause skin irritation. Remove collar at the first sign of irritation or adverse reaction. Collar is intended for use as an insecticide generator and is not to be taken internally by man or animal. Do not use on sick or convalescing animals. DO not administer other cholinesterase inhibitors (i.e. oral pesticides) while collar is worn or for a few days after removing collar.

PRECAUTIONS: Do not open inner envelope until ready to use. DO NOT ALLOW CHILDREN TO PLAY WITH UNATTACHED COLLAR. Dust will form on this collar during storage. Do not get dust or collar in mouth, harmful if swallowed. Wash hands thoroughly with soap and water after handling collar.

FIRST AID: In case of poisoning, call a physician immediately. If dust from collar contacts skin, wash thoroughly with soap and water. If dust contacts eyes, flush eyes with water and get medical attention.

TOXICOLOGICAL INFORMATION: Dust in eyes will cause temporary pupillary constriction\*. Dust released from collar is a cholinesterase inhibitor. Atropine and homatropine\* are antidotal.

DISTRIBUTED BY:  
ROLF C. HAGEN INC.  
3225 SARTELON  
MONTREAL, QUEBEC  
H4R 1E8

\*\*\*\*\*

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))