

BOOT HILL® Rodenticide Mini Block

Kills Rats & Mice Dead

Kills Rats & Mice with a single feeding

Kills Warfarin-Resistant Norway Rats

DOMESTIC

GUARANTEE: Bromadiolone..... 0.005%

REG. NO. 23399 P.C.P. ACT

KEEP OUT OF REACH OF CHILDREN

READ THE LABEL BEFORE USING

WARNING, CONTAINS THE ALLERGENS PEANUTS AND WHEAT

WARNING

{Insert "Diamond" Skull & Crossbones}

POISON

Liphatech Inc.
3600 W. Elm Street
Milwaukee, WI
53209
U.S.A.
(414) 351-1476

Net Contents: 1 Kg

DIRECTIONS FOR USE: Use BOOT HILL® Rodenticide Mini Block for the control of Norway rats, Roof rats and House mice indoors and against the outside walls of buildings.

Selection of Treatment Areas:

USE LIMITATIONS:

Bait MUST either be placed in tamper-resistant bait stations or in locations not accessible to children, pets or livestock. DO NOT place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food.

AREAS OF USE:

Place bait in locations where rats and mice travel and will most likely find and consume the bait. Placement along walls, in corners, by gnawed openings, in and around burrows, and in concealed locations where rats and mice feed is desirable. Maintain a steady supply of fresh bait. Remove any spoiled or contaminated bait promptly.

APPLICATION DIRECTIONS:

For Norway Rats (*Rattus norvegicus*) and Roof Rats (*Rattus rattus*): Norway rats usually inhabit the ground floor of structures and burrow in the ground. They will range approximately 15 to 100 m from their nesting area seeking food and water. Roof rats, however, are excellent climbers and usually infest the upper stories of structures. Hence, special attention must be paid to this species ability to climb by placing baits in the roof rats activity area. For both species, eliminate as much of the naturally occurring food as possible and place 3 to 16 pieces of mini block (usually at intervals of 5 to 10 meters) within the treatment area. Maintain an uninterrupted supply of fresh bait for at least 10 days, or until signs of rat activity cease.

For House Mice (*Mus musculus*): House mice are excellent climbers and will infest all floors in a structure. Mice will only range approximately 10 m from their nesting area seeking food. Eliminate as much of the naturally occurring food as possible and place 1 piece of mini block at intervals of 2 to 4 meters within the treatment area. If a bait placement is not disturbed after several days, move the placement approximately 1 meter. Two (2) pieces of mini block per placement may be necessary at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 days or until signs of mouse activity cease.

Note: Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and unconsumed bait properly. To prevent reinfestation, eliminate rodent food, water and harbourage as much as possible. If reinfestation occurs, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish bait as needed.

PRECAUTIONS:

KEEP OUT OF REACH OF CHILDREN, PETS AND LIVESTOCK. May be harmful or fatal if swallowed or absorbed through the skin. Rubber gloves must be worn when handling this product and when disposing of dead rodents, unconsumed baits and empty containers. Avoid contact with eyes, skin or clothing. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Wash skin thoroughly with soap and water after handling. Wash contaminated clothing, separately from other laundry, with soap and hot water before reuse. KEEP AWAY FROM FEED AND FOODSTUFFS. This product is toxic to fish and wildlife. Keep out of lakes, streams and ponds.

FIRST AID INSTRUCTIONS:

If in eyes, flush with water for 15 minutes. If irritation persists, seek medical attention. If on skin, wash affected area thoroughly with soap and water. If swallowed, give patient one or two glasses of water and induce vomiting by touching back of throat with a finger or blunt object; repeat until vomitus is clear. Do not induce vomiting or give anything by mouth to an unconscious person. Seek medical attention or contact Poison Control Centre immediately.

For all cases of human ingestion, immediately notify a physician or poison control centre. If pet or livestock poisoning is suspected, immediately contact a veterinarian.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION:

Vitamin K1 in the form of intramuscular or subcutaneous injections, or by oral ingestion are suggested remedial treatments for anticoagulant poisoning. The severity of the case measured by establishing prolonged prothrombin times (P.T.) will determine appropriate therapy. Monitoring P.T. will indicate the necessity of repeated treatments.

STORAGE AND DISPOSAL:

Store in a cool, dry place away from other chemicals and food or feed. Store product not in use, in original container, in a secure location inaccessible to children and non-target animals.

Do not contaminate water, food or feed by storage or disposal. Do not reuse empty container. Wrap and dispose of empty container with household garbage.

RODENT DISPOSAL:

Dispose of dead rodents in garbage or by burying.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.