Parapel(R)

DOMESTIC

PARAFFINIZED PELLETS

KILLS NORWAY AND ROOF RATS AND HOUSE MICE. PARAPEL BAITS ARE ACTUALLY IMPREGNATED BY THE TOXICANT.

GUARANTEE:

2-(p-chlorophenyl) 1.3 indandione.....0.005%

REGISTRATION NO. 21371

PEST CONTROL PRODUCTS ACT

READ THE LABEL BEFORE USING

KEEP OUT OF REACH OF CHILDREN

CAUTION

NOTE TO PHYSICIAN: If ingested, administer Vitamin K1, intramuscularly or orally, as indicated in bishydroxycoumarin overdoses. Repeat as necessary based on monitoring of prothrombin times.

The Archem Corp. 1514 Eleventh Street Portsmouth, Ohio 45662 Phone 614-353-1125

NET CONTENTS:

PRECAUTIONS:

KEEP OUT OF REACH OF CHILDREN

Keep away from humans, domestic animals and pets. If swallowed or absorbed by humans, domestic animals or pets, this material may reduce the clotting ability of the blood and cause bleeding.

Do not contaminate water, food or feed by storage or disposal.

ENVIRONMENTAL HAZARD:

Keep Out of Lakes, Streams or Ponds.

DIRECTIONS FOR USE: For control of Norway rats, Roof rats and House mice in and around homes and similar man-made structures.

Treated baits must be placed in tamper proof bait boxes or in locations not accessible to children, pets, domestic animals or wildlife. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food.

SELECTION OF TREATMENT AREAS

Determine area where rats and/or mice will most likely find and consume the bait. Generally, these areas are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been observed. Remove as much food as possible.

APPLICATION DIRECTIONS

RATS: Apply 112 to 448 grams of bait per placement. Maintain an uninterrupted supply of fresh bait for at least 10 days.

MICE: Apply 7 to 14 grams (1-2 level tablespoons) of bait at 8 to 12 foot intervals. Larger placements (up to 56 grams) may be need at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 days.

RATS AND MICE: Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and unconsumed bait properly. To prevent reinfestation, eliminate food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish bait as needed.

FIRST AID

IF SWALLOWED: Call a physician or Poison Control Centre. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

TOXICOLOGY: If swallowed by humans, domestic animals, or pets,

this material may reduce the clotting ability of the blood and cause bleeding. In such case intramuscularly and oral administration of Vitamin K1 combined with blood transfusions are indicated as in the case of hemorrhage caused by overdoses of bishydroxycoumarin.

STORAGE: Store only in original container, in a cool, dry place inaccessible to children and pets.

DISPOSAL: Do not reuse container. Wrap and dispose of with regular household garbage.

PARAPEL BAIT IS OIL IMPREGNATED!

To ensure the most effective and highest quality bait, this product has been prepared in a two-step operation.

- 1. The active ingredient was dissolved in mineral oil to form a concentrate.
- 2. This mineral oil concentrate was then used to impregnate the bait.

This process give the following advantages:

- a. The mineral oil aids in the preservation of the bait, thus reducing the possibility of rancidity that may cause bait rejection.
- b. Impregnation reduces the possibility of the toxicant detaching itself from the bait.

Seller's guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

Packed by Batch No.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))