

2001/2002

CO-OP (R)

Bug Killer

For control of vegetable and fruit insects

Insecticide Dust

DOMESTIC

READ THE LABEL BEFORE USING

GUARANTEE: Carbaryl 5%

REGISTRATION NO. 10645
PEST CONTROL PRODUCTS ACT

2 kg/ 500 g

Interprovincial Cooperative Limited
P. O. BOX 1050, Saskatoon, Saskatchewan S7K 3M9
Winnipeg Mississauga Montreal Moncton

DIRECTIONS: For small areas, apply a light even coating of dust to the upper and lower leaf surfaces. For larger areas, apply the rate indicated below. Use lower rates on young plants.

	g Dust per 100 m²	Remarks
Potatoes, Tomatoes, Eggplant, Pepper: Colorado Potato Beetle, Potato Flea Beetle, Leaf Hoppers, Fall Army Worm, Tomato Horn Worm,	250 - 600 g	Apply when insects or damage first appears. Repeat in 7-10 days or as necessary. Allow 7 day interval between last dust and harvest.
Lygus Bug, Stink Bug.	600 g	

Cucumbers, Squash, Melons, Pumpkins: Cucumber Beetles, Squash Bugs, Flea Beetles, Leaf Hoppers.	300 - 500 g	Some leaf injury may occur if treatments are made when young foliage is wet or during prolonged high humidity. Allow 3 day interval between last dust and harvest.
Corn: European Corn Borer	250 - 500 g	Make 4 or more application 5 days apart. Start when eggs are laid. Allow 1 day interval between last dust and harvest.
Corn: Earworm	250 - 500 g	Make 4 applications 2-4 days apart. Start at 25% silk showing.
Fall Army Worm	250 - 500 g	
Beans: Mexican Bean Beetles	300 g	Apply when insects or damage appears. Repeat at 7-10 day intervals. Allow 3 day interval between last dust and harvest.
Leaf Hopper, Flea Beetle	350 g	
Lygus and Stink Bug	500 g	
Strawberries: Meadow Spittle Bug Strawberry Leaf Roller.	350 - 500 g	Apply when insects or damage appears. Repeat at 7-10 day intervals or as necessary. Allow 1 day between last dust and harvest.
Blueberries: Blueberry Maggot, Cherry and Cranberry Fruit Worm, Japanese Beetle.	200 g	Apply three weeks before harvest and repeat 10 days later. Allow 1 day interval between last dust and harvest.

Grapes: Grape Berry Moth, Leaf Hoppers.	400 - 600 g	Apply in morning or evening when vines are damp. Allow 3 day interval between last dust and harvest.
--	--------------------	--

PRECAUTIONS: KEEP OUT OF THE REACH OF CHILDREN. Harmful if swallowed or inhaled. Avoid breathing dust. Avoid contact with skin. Wash thoroughly after handling. Change contaminated clothing. Toxic to bees. Do not apply during pollination. Do not contaminate streams, ponds and water supplies.

FIRST AID:

1. In case of poisoning call a physician or contact a poison control centre immediately.
2. If on skin, remove contaminated clothing and wash skin thoroughly with soap and water.
3. If in eyes, flush with water immediately and continue for 5-10 minutes. Get medical attention or contact a poison control centre.
4. If swallowed, induce vomiting then take patient to nearest hospital or doctor's office or contact a poison control centre. Take the pesticide label with you.

TOXICOLOGICAL INFORMATION: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal. Do not use 2-PAM, opiates or cholinesterase inhibiting drugs.

DISPOSAL: Do not reuse empty container. Wrap empty container and discard in household garbage.

NOTICE TO BUYER: Seller's guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

(R)Registered Trademark of TMC Distributing Ltd.
Used under license

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from

reliance upon this service.

+))