

96.03.05

CO-OP®

Aero-Tack

Agricultural Insecticide
Pressurized Spray

A concentrated insecticide for use in
Milk Parlours and Milkhouses and on
Dairy and Beef Cattle and Horses to
kill flying insects.

Also for use in Agricultural and other
Industrial Buildings, to kill flying and
crawling insects.

AGRICULTURAL

GUARANTEE:

Pyrethrins 0.50%
Piperonyl Butoxide 4.00%

REGISTRATION NO. 24413

PEST CONTROL PRODUCTS ACT

READ THE LABEL BEFORE USING

DANGER

EXTREMELY FLAMMABLE EXPLOSIVE

600 g

Interprovincial Cooperatives Limited
P.O. Box 1050
Saskatoon, Saskatchewan
S7K 3M9

WINNIPEG MISSISSAUGA MONTREAL MONCTON

DIRECTIONS: Hold can upright while spraying. SHAKE WELL BEFORE USING.

FLYING INSECTS: Flies, Mosquitoes, Wasps, and Flying Moths. Close doors and windows. Apply a 6-10 second burst per 30m³ of space. Direct mist upwards in sweeping motion to all parts of the room, especially windows and other light sources. Also direct mist into areas where insects are present. Insects must be contacted to be killed. Keep spray at least 1 m from interior walls, fabrics and furniture. Keep doors and windows closed for 15 minutes after each treatment. Ensure proper ventilation after sealed areas have been treated. Clean up dead insects. Spray only when temperature is under 32EC.

CRAWLING INSECTS: Cockroaches, spiders, crickets, ants, carpet beetles, centipedes and silver fish. Spray hiding places such as baseboards, moist areas, opening around sinks, drains and pipes, behind cabinets and storage areas. Spray ant trails, around door frames, window sills, or in ant nests. Insects must be contacted with spray to be killed.

FLEAS: Spray infested areas such as pet's bed and resting quarters, nearby cracks and crevices, along and behind baseboards, window and door frames, and localized areas of floor and floor covering where these pests may be present. Remove old bedding and replace with fresh, clean bedding after treatment. Insects must be contacted to be killed. DO NOT TREAT PETS WITH THIS PRODUCT.

SPACE SPRAY: Kills flying insects in Milking Parlours and Milking Rooms. Direct spray upwards to all parts of the room especially windows and other light sources. Apply at a rate of 6 to 10 seconds per 30m³. Remove all animals from treatment area before using a space spray. For best results, close all doors and windows during treatment and keep room closed for fifteen minutes before ventilating. Ventilate areas thoroughly for at least 30 minutes before animal re-entry. Insects must be contacted to be killed. Sweep up dead insects. Repeat as necessary with additional spraying of breeding areas. Spray only when temperature is under 32EC.

LIVESTOCK SPRAY: To treat Beef, Dairy Cattle and Horses against attack of houseflies, stable flies and horn flies, spray animals once daily with a fine mist. Apply at a rate of 6 to 8 seconds per animal. Spray should be enough to wet ends of the hair only. Do not wet the hair or skin. Take care not to spray in animal's eyes or mouth. Treat dairy cattle at least 20 minutes before milking. Do not spray directly on udder. Wash udders before milking. Do not use more often than once a day.

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN. Do not remain in treated areas. Ensure proper ventilation after sealed areas have been treated. Cover milk and milking utensils to prevent contamination from spray and dead insects. Do not contaminate feed or water troughs. Do not spray over milking equipment. (Any treated surfaces that may contact food or feed products should be washed thoroughly with potable water after spraying and before re-use). Avoid contact with skin and eyes. Wash after using. CONTENTS UNDER PRESSURE: Do not place in hot water or near radiators, stoves or other sources of heat. Do not puncture or incinerate can or store at temperature over 50°C. Do not spray into or near any open flame. Not for use on humans. Avoid inhalation of spray.

DO NOT USE IN THE PRESENCE OF AN OPEN FLAME OR SPARK.

FIRST AID: If sprayed in eyes or on skin, flush with clean water. If irritation persists, call a Poison Control Centre and obtain medical attention. If accidentally swallowed, obtain medical attention at once.

DISPOSAL: When container is empty, press valve to release all remaining pressure. Dispose of empty container with household garbage. For information on the disposal of unused, unwanted product and the cleanup of spills, contact the Provincial Regulatory Agency or the Manufacturer.

NOTICE TO USER:

This control product is to be used only in accordance with the directions on this label. It is an offense under the Pest Control Products Act to use a control product under unsafe conditions.

LIMITATIONS OF WARRANTY:

Seller's guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any

loss resulting, directly or indirectly, from reliance upon this service.

+))