DISVAP AQUA

BARN AND LIVESTOCK INSECTICIDE WATER BASE MICRO-EMULSION

READ THE LABEL BEFORE USING

COMMERCIAL

GUARANTEE (W/V)

To control stable flies, house flies, horseflies, mosquitoes and gnats.

KEEP OUT OF REACH OF CHILDREN

REGISTRATION NO. 22202

PEST CONTROL PRODUCTS ACT

NET CONTENTS: 4 L

DISPAR, DIVISION OF VETOQUINOL N.-A. Inc. Lavaltrie, Qc, Canada JOK 1H0

INSECTICIDE DISVAP AQUA

DIRECTIONS: LIVESTOCK SPRAY: To control stable flies, house flies, horse flies, mosquitoes and gnats: apply 30-60 mL per adult animal to wet the hair thoroughly paying particular attention to legs, flanks, back, trunk and other body areas commonly attacked by these flies. Repeat weekly or as needed. To control face flies, horn flies, horse flies on beef cattle, dairy cattle (lactating or not) hogs, horses. Apply to the face of the animal in the morning before releasing to pasture. Apply sufficiently to wet the face, but no more than 30 to 45 mL per animal. DO NOT spray in animal's eyes or mouth. Repeat daily as needed. To kill fleas on horses, cattle and goats, and to protect against reinfestation, wet the animal by spraying. Repeat as necessary.

Treat dairy cattle at least 20 minutes before milking. DO NOT spray directly on udders. Wash udders before milking. DO NOT cover horses after treating.

SPACE SPRAY: In hog, dairy and beef barns, milking parlours, milk rooms, dairies and poultry houses. To control house flies, stable flies, mosquitoes and gnats. Apply as a fog or fine mist, directing spray towards the ceiling and upper corners until area is filled with mist, using about 60 mL per 30 m³ of space. For best results, close doors and windows before spraying, and keep them closed for 10 to 15 minutes. Vacate the area and ventilate before reentry. Repeat application as necessary.

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN. Not for use on humans. Harmful if swallowed. Avoid contact with skin, eyes and clothing, and avoid inhalation of spray. Wear mask or respirator while fogging. Keep container closed. Wash after use. Do not remain in treated areas. Ensure proper ventilation after sealed areas have been treated. Any treated surface that may contact food or feed products should be washed thoroughly with potable water after spraying and before use. Cover milk and milk utensils to prevent contamination from spray and dead insects. Do not use in areas where beekeeping equipment or supplies are stored. Product toxic to fish. Do not contaminate any body of water.

DISPOSAL: Rinse empty container thoroughly and add rinse water to fogging machine tank. Follow Provincial instructions for any required cleaning of container prior to its disposal. Dispose of empty container in accordance with provincial requirements. For more information on the disposal of unused or unwanted products and the clean up of spills, contact the provincial regulatory agency or the manufacturer.

FIRST AID: If in eyes or on skin, flush with clean water, if irritation persists, obtain medical attention. If accidentally swallowed, get prompt medical attention.

NOTICE TO USER: This control product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products Act to use a control product under unsafe conditions.

LIMITATION OF WARRANTY: Seller's guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risks to persons or property arising from the use or handling of this product and accepts the product on that condition.

STORAGE: Do not expose to temperatures below OEC.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))