

85.12.31

BAYTEX (R)

1% GRANULAR

RESIDUAL MOSQUITO LARVICIDE

Contains fenthion

RESTRICTED

NET WEIGHT _____ KILOGRAMS

GUARANTEE: Fenthion.....1%

REGISTRATION NO. 9676 PEST CONTROL PRODUCTS ACT

CAUTION

KEEP OUT OF REACH OF CHILDREN

(See rear panel for first aid and caution statements)

READ THE LABEL BEFORE USING

CHEMAGRO LIMITED
1355 Aerowod Drive
Mississauga, Ontario, Canada L4W 1C2

NOTICE TO USER: This control product is to be used only in accordance with the directions on this label. It is an offence under the PEST CONTROL PRODUCTS ACT to use a control product under unsafe conditions.

NATURE OF RESTRICTION: This product is to be used only in the manner

authorized; consult local pesticide regulatory authorities regarding appropriate use permits which may be required.

DIRECTIONS FOR USE

BAYTEX 1% Granular mosquito larvicide is recommended for use wherever mosquitoes breed. It provides residual control and is effective against DDT-resistant as well as non-resistant strains.

For mosquito larvae control, applications should be made to moist areas, standing water, shallow ponds, and around the edges of larger ponds, lakes, marshes, swamps, and tidal areas. Also apply around and in dumps, catch basins, and similar areas where mosquitoes breed.

RECOMMENDED APPLICATIONS

INSECT	DOSAGE BAYTEX 1% GRANULAR	REMARKS
Mosquito Larvae	55 to 110 g/100m ² of water surface or 5.5 to 11 kg/ha of water surface	Apply specified dosage uniformly. Use the higher rate where pools of water are greater than 30 cm in depth. Allow 21 days between applications.

PRECAUTIONS

1. Do not use inside homes, in dairy barns, poultry houses, or buildings where food is processed.
2. Do not use on crops used for food.
3. Birds feeding in treated areas may be harmed at the maximum recommended mosquito control rate.

CAUTION

Harmful if swallowed, inhaled, or absorbed through the skin. Do not get in eyes, on skin, or on clothing. Do not breathe dust. Wash thoroughly with soap and warm water after handling. Wash clothing with soap and hot water before reuse.

Avoid contamination of feed or food. Keep out of reach of children.

FIRST AID: In case of poisoning call a physician immediately. Have patient lie down and keep quiet. If swallowed, vomiting should be induced. Administer milk or water freely and induce vomiting by giving one dose (15 mL) of syrup of ipecac. If vomiting does not occur within 10 to 20 minutes, administer second dose. If syrup of ipecac is not available, induce vomiting by sticking finger down throat. Repeat until vomit fluid is clear. The patient should be lying down with the head below the foot level and facing down or to one side. Professional medical assistance should be secured immediately. DO NOT INDUCE VOMITING TO AN UNCONSCIOUS PERSON OR TO PERSONS IN A CONVULSIVE STATE. If on skin, remove contaminated clothing and wash skin immediately with soap and warm water. If eyes are contaminated, wash immediately with flowing water for at least 15 minutes.

TOXICOLOGICAL INFORMATION: The compound inhibits cholinesterase resulting in stimulation of the central nervous system. This results in a sense of "tightness" in the chest, sweating, contracted pupils, stomach pains, vomiting and diarrhea. The antidote is atropine sulfate administered in large therapeutic doses repeated as necessary to the point of tolerance. 2-PAM is also antidotal and may be administered in conjunction with atropine. Do not give morphine. Watch for pulmonary edema, which may develop in serious cases of poisoning even after 12 hours. At first sign of pulmonary edema, the patient should be placed in an oxygen tent and treated symptomatically.

NOTICE

Seller's guarantee shall be limited to the terms set out on the label and, subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

BAYTEX is a Reg. TM of Bayer AG.

Printed in U.S.A.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon

this service.

+))