

19-JAN-2001

X-CELL 425LC

COMMERCIAL

GUARANTEE: Dazomet (3,5-dimethyl tetrahydro-2H-1,3,5-thiadiazine-2-thione)-20%

REGISTRATION NO.: 26664
PEST CONTROL PRODUCTS ACT

WARNING

POISON

READ THE LABEL BEFORE USING

JOHNSON DIVERSEY CANADA INC.

~~DIVERSEY LEVER CANADA~~

2401 Bristol Circle
Oakville, Ontario
L6H 6P1

Notification Change

Emergency Number: (905)-829-1200

Net Contents - 1,045 Kg.

Diversey Lever

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN

WARNING! Causes Eye and Skin Irritation.
Potential Skin Sensitizer.
Harmful if Swallowed or Inhaled.

- ⊘ WEAR CHEMICAL RESISTANT GLOVES WHEN HANDLING THIS PRODUCT.
- ⊘ AVOID CONTACT WITH EYES, SKIN AND CLOTHING.
- ⊘ WEAR GOGGLES OR FACE SHIELD.
- ⊘ DO NOT BREATHE VAPOURS.
- ⊘ USE WITH ADEQUATE VENTILATION.
- ⊘ DO NOT SPRAY THE PRODUCT IN OPEN SYSTEMS.

- ⊃ KEEP CONTAINER CLOSED.
- ⊃ WASH THOROUGHLY AFTER HANDLING.
- ⊃ AVOID CONTAMINATION OF FOOD, FEED OR DRINKING WATER.
- ⊃ LAUNDRER PROTECTIVE CLOTHING BEFORE REUSE.

TOXICOLOGICAL INFORMATION: There is not specific antidote. Medical personnel should provide symptomatic or supportive treatment. Probable mucosal damage may contraindicate the use of gastric lavage.

ENVIRONMENTAL HAZARD: This pesticide is toxic to fish. Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, or public waters. Open dumping is prohibited.

NOTICE TO USER: This control product is to be used only in accordance with the directions on this label. It is an offense under the Pest Control Products Act to use a control product under unsafe conditions.

NOTICE TO BUYER: Seller's Guarantee shall be limited to the terms set out on the label and, subject thereto, the buyer assumes the risk to persons or property arising from the use of handling of this product and accepts the product on that condition.

STORAGE

Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited. If plastic, do not reuse empty container.

DISPOSAL

1. Close Valve.
2. For disposal, this container is to be returned to point of purchase (distributor/dealer). It must be refilled by the distributor with the same product. Do not reuse this container for any other purpose.
3. Contact the distributor/dealer to arrange for the return of the container.
3. For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency.
4. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

DO NOT FLUSH INTO SURFACE STREAMS. INFORM THE PROVINCIAL REGULATORY AGENCY OR THE MANUFACTURER AS IN NUMBER 3.

DIRECTIONS FOR USE

FOR THE PRESERVATION OF CLAY SLURRIES, ADHESIVES, COATINGS AND HIGH VISCOSITY SUSPENSIONS: For preservation of slurries and high viscosity suspensions, X-

CELL 425LC should be added at a point in the processing system where there will be sufficient time and agitation for good mixing and dispersion. Add X-CELL 425LC at use levels of 0.04-0.40% by weight, based on the total formulation in slurries of starch, clay, calcium carbonate or titanium dioxide; paper coatings; high viscosity suspensions (e.g. polymers; silica-polymer combinations); polyvinyl alcohol/polyvinyl acetate based adhesives; starch based adhesives; dextrin based adhesives. The exact amount of material to be added for the preservation of any given formulation will depend on the components and local storage conditions. Dosage rates should be determined by actual tests.

PAPER MILL SLIME CONTROL: FOR THE CONTROL OF SLIME-FORMING AND/OR SPOILAGE BACTERIA:

X-CELL 425LC is added at a point in the system (raw stock chest; beater and/or refiner chest or machine chest-wire pit) where it will be uniformly mixed. Application may be continuous or intermittent for a certain number of hours/day or per shift, depending upon system characteristics.

Add 160 to 650 ml of X-CELL 425LC per metric ton of paper produced.

INTERMITTENT FEED METHOD: Apply 390 to 650 ml of X-CELL 425LC per metric ton (dry basis) of pulp or paper for 2 hours every 8 hours. Badly fouled process systems must be cleaned before initial treatment.

CONTINUOUS FEED METHOD: Apply 160 to 480 ml of X-CELL 425LC per metric ton (dry basis) of pulp or paper produced on a continuous basis. Badly fouled process systems must be cleaned before initial treatment.

FIRST AID

In case of contact, immediately flush eyes and skin with large amounts of water for at least 15 minutes. Remove contaminated clothing and shoes. For eyes, hold eyelids apart during the flushing to ensure rinsing of the entire surface of the eye and lids with water. Get medical attention. Wash clothing before reuse.

If swallowed, immediately call a physician, a poison control centre, or the nearest hospital. Induce vomiting as directed by medical personnel. Never give anything by mouth to an unconscious or convulsing person. Get medical attention immediately. Take container, label or product name and Pest Control Product Registration number with you when seeking medical attention.

If inhaled, remove to fresh air. If not breathing, give artificial respiration. If breathing is difficult, give oxygen. Get medical attention immediately

LABEL DATE: 09-Nov-00

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through

this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))