

97.04.22

(( INSERT ))

PRO-SPOT™ 20

Solution for Flea Control on Dogs

FOR VETERINARY USE ONLY

COMMERCIAL

CAUTION POISON

GUARANTEE:

fenthion .....5.6%

REGISTRATION NO. 22033 PEST CONTROL PRODUCTS ACT

KEEP OUT OF REACH OF CHILDREN

READ THE LABEL BEFORE USING

NET CONTENTS:

50 x 0.74 mL Tubes

Bayer Inc.  
77 Belfield Road  
Etobicoke, Ontario  
M9W 1G6

NOTICE TO USER: This control product is to be used only in accordance with the directions on this label. It is an offense under the PEST CONTROL PRODUCTS ACT to use a control product under unsafe conditions.

DIRECTIONS FOR USE

Name	Dog Weight	Concentration	mg a.i./m L	Net Contents
Pro-Spot™ 20	5.0-9.1 kg	5.6%	54	0.74 mL

Dogs should be weighed prior to selection and use of Pro-Spot™ to ensure the recommended dose is applied. Dogs with body weights of less than 5.0 kg should not be treated with this product. Apply the contents of one single-dose tube to the dog. Young, growing dogs should be reweighed periodically to ensure the proper tube of Pro-Spot™ is used for repeat treatments. The frequency of repeat treatments depends on the rate of flea reinfestation. This will vary greatly with season and location. Treatments should not be repeated more often than once every 2 weeks.

#### HOW TO APPLY:

1. Wear chemical-resistant gloves during application. Hold the applicator tube in an upright position. Pull cap off tube then turn cap around and place other end of cap back on tube. Twist cap to break seal, then remove cap from tube.
2. Invert the tube and use it to part the hair on the dog's back between the shoulder blades.
3. Squeeze the tube firmly twice to apply all the solution in one spot directly on the skin. While applying the liquid, the tube should remain in contact with the dog's skin to ensure that the total dose is properly applied. Do not apply on top of hair as this will result in reduced flea control. Care should be taken to avoid contact of solution with human skin.
4. Wrap empty tube in paper and discard in garbage.

For effective flea control, treatment of the pet should be combined with the sanitation of any area used by the pet. Vacuum floors, carpets and furniture (discard vacuum bag after use) and wash the pet's bedding, living quarters and surrounding areas. If pest problems persist, an insecticidal premise treatment may be required.

#### USE LIMITATIONS:

Fenthion is a cholinesterase-inhibitor. Do not use this product on dogs at the same time or within 10 days before or after treatment with other cholinesterase inhibiting pesticides, drugs or chemicals. Do not use with flea or tick collars. Do not use more often than once every 2 weeks. Do not use on puppies under 10 weeks of age. Do not use on sick or convalescing dogs. The safe use of Fenthion in breeding males and pregnant females has not been established.

Treatment with Pro-Spot™ at 2-week intervals should not exceed 6

months as controlled safety studies of longer duration have not been conducted.

POSSIBLE ADVERSE REACTIONS:

May cause irritation of the skin at the application site. Some animals may lick and scratch the application site, and may require control to prevent self-mutilation. Animals that have been previously exposed to organophosphate ester insecticides are at greater risk of an adverse reaction following exposure to Pro-Spot™.

OVERDOSAGE:

Acute symptoms of overdose in dogs include diarrhea, salivation, vomiting, muscular tremors and weakness. Consult veterinarian at first sign of adverse effects.

NOTICE TO VETERINARIAN:

If toxicity should occur, atropine is antidotal.

PRECAUTIONS:

KEEP OUT OF REACH OF CHILDREN. May be harmful if swallowed or absorbed through the skin. Do not get in eyes, on skin or on clothing. Avoid breathing vapours. Wear chemical resistant gloves when applying this product. Wash thoroughly with soap and warm water immediately after handling. Wash contaminated clothing with soap and hot water before re-use. Do not contaminate feed or food.

This material contains a cholinesterase inhibitor.

STORAGE:

Keep single dose applicators stored in their original container until use. Do not store near heat or open flame.

DISPOSAL:

Wrap empty tubes in paper and discard in garbage.

FIRST AID:

If swallowed, contact physician or poison control centre immediately. If spilled on skin, wash immediately with soap and warm water. If in eyes, rinse immediately with large amounts of water and obtain prompt medical aid.

TOXICOLOGY INFORMATION:

Atropine is antidotal. 2-PAM is also antidotal, but must be administered in conjunction with atropine.

NOTICE:

Seller's guarantee shall be limited to the terms set out on the label and, subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

(( LABEL ))

PRO-SPOT™ 20

Solution for Flea Control on Dogs

FOR VETERINARY USE ONLY

COMMERCIAL

GUARANTEE:

fenthion .....5.6%

REGISTRATION NO. 22033 PEST CONTROL PRODUCTS ACT

NET CONTENTS

50 X 0.74 mL Tubes

CAUTION POISON

KEEP OUT OF REACH OF CHILDREN

READ THE LABEL BEFORE USING

Bayer Inc.  
77 Belfield Road  
Etobicoke, Ontario  
M9W 1G6

NOTICE TO USER: This control product is to be used only in

accordance with the directions on this label. It is an offense under the PEST CONTROL PRODUCTS ACT to use a control product under unsafe conditions.

#### DIRECTIONS FOR USE

Name	Dog Weight	Concentration	mg a.i./m L	Net Contents
Pro-Spot™ 20	5.0-9.1 kg	5.6%	54	0.74 mL

Dogs should be weighed prior to selection and use of Pro-Spot™ to ensure the recommended dose is applied. Dogs with body weights of less than 5.0 kg should not be treated with this product. Apply the contents of one single-dose tube to the dog. Young, growing dogs should be reweighed periodically to ensure the proper tube of Pro-Spot™ is used for repeat treatments. The frequency of repeat treatments depends on the rate of flea reinfestation. This will vary greatly with season and location. Treatments should not be repeated more often than once every 2 weeks.

#### HOW TO APPLY:

1. Wear chemical-resistant gloves during application. Hold the applicator tube in an upright position. Pull cap off tube then turn cap around and place other end of cap back on tube. Twist cap to break seal, then remove cap from tube.
2. Invert the tube and use it to part the hair on the dog's back between the shoulder blades.
3. Squeeze the tube firmly twice to apply all the solution in one spot directly on the skin. While applying the liquid, the tube should remain in contact with the dog's skin to ensure that the total dose is properly applied. Do not apply on top of hair as this will result in reduced flea control. Care should be taken to avoid contact of solution with human skin.
4. Wrap empty tube in paper and discard in garbage.

For effective flea control, treatment of the pet should be combined with the sanitation of any area used by the pet. Vacuum floors, carpets and furniture (discard vacuum bag after use) and wash the pet's bedding, living quarters and surrounding areas. If pest problems persist, an insecticidal premise treatment may

be required.

USE LIMITATIONS:

Fenthion is a cholinesterase-inhibitor. Do not use this product on dogs at the same time or within 10 days before or after treatment with other cholinesterase inhibiting pesticides, drugs or chemicals. Do not use with flea or tick collars. Do not use more often than once every 2 weeks. Do not use on puppies under 10 weeks of age. Do not use on sick or convalescing dogs. The safe use of Fenthion in breeding males and pregnant females has not been established.

Treatment with Pro-Spot™ at 2-week intervals should not exceed 6 months as controlled safety studies of longer duration have not been conducted.

POSSIBLE ADVERSE REACTIONS:

May cause irritation of the skin at the application site. Some animals may lick and scratch the application site, and may require control to prevent self-mutilation. Animals that have been previously exposed to organophosphate ester insecticides are at greater risk of an adverse reaction following exposure to Pro-Spot.

OVERDOSAGE:

Acute symptoms of overdosage in dogs include diarrhea, salivation, vomiting, muscular tremors and weakness. Consult veterinarian at first sign of adverse effects.

NOTICE TO VETERINARIAN:

If toxicity should occur, atropine is antidotal.

PRECAUTIONS:

KEEP OUT OF REACH OF CHILDREN. May be harmful if swallowed or absorbed through the skin. Do not get in eyes, on skin or on clothing. Avoid breathing vapours. Wear chemical resistant gloves when applying this product. Wash thoroughly with soap and warm water immediately after handling. Wash contaminated clothing with soap and hot water before re-use. Do not contaminate feed or food.

This material contains a cholinesterase inhibitor.

STORAGE:

Keep single dose applicators stored in their original container until use. Do not store near heat or open flame.

DISPOSAL:

Wrap empty tubes in paper and discard in garbage.

FIRST AID:

If swallowed, contact physician or poison control centre immediately. If spilled on skin, wash immediately with soap and warm water. If in eyes, rinse immediately with large amounts of water and obtain prompt medical aid.

TOXICOLOGY INFORMATION:

Atropine is antidotal. 2-PAM is also antidotal, but must be administered in conjunction with atropine.

NOTICE:

Seller's guarantee shall be limited to the terms set out on the label and, subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

(( APPLICATOR TUBE ))

PRO-SPOT™ 20

READ THE LABEL AND INSERT BEFORE USING

GUARANTEE:  
Fenthion 5.6%

REGISTRATION NO. 22033 P.C.P. ACT

CAUTION POISON

BAYER INC.  
1-800-268-1331

\*\*\*\*\*

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label

information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))