

96.07.23

0910-0911

ANTISTAF

DIN 656224

DISINFECTANT-CLEANER-SANITIZER-FUNGICIDE

GUARANTEE

Octyl Decyl Dimethyl Ammonium Chloride.....1.14%
Dioctyl Dimethyl Ammonium Chloride.....0.57%
Didecyl Dimethyl Ammonium Chloride.....0.57%
Alkyl (C14,50%; C12,40%; C16, 10%),
Dimethyl Benzyl Ammonium chloride.....1.52%

MILDEWSTAT-VIRUCIDE-DEODORIZER
WITH ORGANIC SOIL TOLERANCE
FOR HOSPITALS, INSTITUTIONAL AND INDUSTRIAL USE

REGISTRATION NO. 19921
PEST CONTROL PRODUCTS ACT

READ THE LABEL BEFORE USING

FOR INDUSTRIAL OR INSTITUTIONAL USE ONLY

NOT FOR RETAIL SALE

CAUTION

POISON\CORROSIVE

BRISK

NET CONTENTS: 4 L 4 X 4 L 20 L

BRISK CORPORATION
Head Office:
480 boul. Montpellier, St-Laurent, Quebec. H4N 2G7

KEEP OUT OF REACH OF CHILDREN

ANTISTAF has been formulated to aid in the reduction of cross-contamination in institutions, schools and hospitals. ANTISTAF is a one-step disinfectant-cleaner that is effective against a broad spectrum of bacteria, is virucidal, fungicidal, and will prevent the growth of mold and mildew and their odors when used as directed.

ANTISTAF is an effective one-step sanitizer-cleaner for use on non-food contact surfaces. When used as directed, ANTISTAF will deodorize surfaces in toilet areas, behind and under sinks and counters, garbage cans and garbage storage areas, and other places where bacterial growth can cause malodorous.

PRECAUTIONARY STATEMENTS: Danger: corrosive to eyes. Avoid contact with eyes, skin and clothing. Harmful if swallowed. Avoid contamination of food.

DISPOSAL: 1. Rinse emptied container thoroughly and add the rinsings to the treatment site. 2. Follow provincial instructions for any required additional cleaning of the container prior to its disposal. 3. Make the empty container unsuitable for further use. 4. Dispose of the container in accordance with provincial requirements. 5. For information on the disposal of unused, unwanted product and the cleanup of spills, contact the Provincial Regulatory Agency or the Manufacturer.

FIRST-AID: In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, CALL A PHYSICIAN. Remove contaminated clothing under running water. INGESTION: Drink promptly a large quantity of milk, egg whites, gelatin solution; or if these are not available, drink large quantities of water. Avoid alcohol. CALL A PHYSICIAN OR POISON CONTROL CENTRE IMMEDIATELY.

TOXICOLOGICAL INFORMATION: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression, and convulsion may be needed.

NOTICE TO USER: This control product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products Act to use a control product under unsafe conditions.

USE DIRECTIONS

ANTISTAF is a proven "one-step" cleaner-disinfectant-sanitizer-fungicide-mildewstat-virucide in the presence of moderate amounts of organic soil.

Apply ANTISTAF to walls, floors and other hard (inanimate) non-porous surfaces such as tables, chairs, countertops, sinks and bedframes, with a cloth, mop or mechanical spray device so as to thoroughly wet surfaces. For heavily soiled areas a preliminary cleaning is required. Prepare a fresh solution daily or when use solution becomes visibly dirty.

DISINFECTION - To disinfect hard non-porous surfaces in hospitals, add: 17 ml per litre of water. Treated surfaces must remain wet for 10 minutes. At this use-level ANTISTAF is also virucidal, fungicidal and mildewstatic at 9 mL per litre of water.

ANTISTAF will disinfect non-porous surfaces in schools, industry and non-medical institutions.

SANITIZING - To sanitize, non-porous non-food contact surfaces, add 5 ml per litre of water. Treated surfaces must remain wet for 60 seconds.

FOR HOSPITALS AND NURSING HOMES: Add 17 ml per litre of water. The broad spectrum of ANTISTAF has been evaluated in the presence of 5% serum and found to be effective against the following organisms by the AOAC Use-Dilution test.

Pseudomonas aeruginosa	Staphylococcus aureus
Salmonella choleraesuis	Escherichia coli
Serratia marcescens	Klebsiella pneumoniae
Enterobacter aerogenes	Salmonella schottmuelleri
Streptococcus faecalis	Shigella dysenteriae
Brevibacterium ammoniagenes	

AOAC Fungicidal test

ANTISTAF at 9 ml per litre of water is an effective fungicide against Trichophyton mentagrophytes (the athlete's foot fungus) when used on surfaces in areas such as locker rooms, dressing rooms, shower and bath areas, exercise facilities, etc.

MOLD AND MILDEW CONTROL: At 9 ml per litre of water ANTISTAF will effectively inhibit the growth of mold and mildew and the odors caused by them when applied to hard, non-porous surfaces (as indicated in general instructions above). Allow to dry on surface and repeat when mildew growth returns.

VIRUCIDAL PERFORMANCE: At 9 ml per litre of water use-level ANTISTAF was evaluated in the presence of 5% serum and found to be effective against influenza A/Brazil Virus on inanimate environmental surfaces.

FOR SCHOOLS, NON-MEDICAL INSTITUTIONAL AND INDUSTRIAL USES: At 9 ml per litre of water ANTISTAF delivers excellent cleaning and germicidal effectiveness. It is effective against Staphylococcus

aureus, Salmonella choleraesuis and Escherichia Coli. The same AOAC tests used above to confirm suitable performance for hospitals were used.

SANITIZING NON-FOOD CONTACT SURFACES: (such as floors, walls, tables, etc). At 5 ml per litre of water use-level ANTISTAF is an effective sanitizer against Staphylococcus aureus and Klebsiella pneumoniae in the presence of 5% serum on non-porous environmental surfaces.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))