2000.03.14

3M

QUAT DISINFECTANT CLEANER CONCENTRATE

WARNING

POISON

CORROSIVE

KEEP OUT OF REACH OF CHILDREN

READ THE LABEL BEFORE USING

REGISTRATION NO. 23652 PEST CONTROL PRODUCTS ACT

DIN 02091852

GUARANTEE COMMERCIAL:

OCTYL DECYL DIMETHYL AMMONIUM CHLORIDE	6.510%
DIOCTYL DIMETHYL AMMONIUM CHLORIDE	2.604%
DIDECYL DIMETHYL AMMONIUM CHLORIDE	3.906%
ALKYL (C12 40%; C14 50%; C16 10%)	
DIMETHYL BENZYL AMMONIUM CHLORIDE	8.680%

2 LITRES

COMMERCIAL CARE DIVISION IMPORTED BY 3M CANADA INC., POST OFFICE BOX 5757 5757, LONDON, ONTARIO N6A 4T1 MADE IN U.S.A.

3M Quat Disinfectant Cleaner Concentrate

For use in institutions, schools, hospitals and food processing plants.

Directions For Use

3M Quat Disinfectant Cleaner is a proven "one-step" disinfectantcleaner-sanitizer-fungicide-mildewstat-virucide which is effective in water up to 400 pm hardness in the presence of 5% serum contamination. Apply 3M Quat Disinfectant Cleaner to walls, floors and other hard (inanimate) non-porous surfaces such as tables, chairs, countertops, sinks, tile, porcelain and bedframes with a cloth, mop or trigger spray device so as to thoroughly wet surfaces. For heavily soiled areas, a preliminary cleaning is required. Prepare a fresh solution daily or when use-solution becomes visibly dirty. All surfaces which come in contact with food are to be rinsed after disinfection and prior to use.

DISINFECTION: To disinfect, hard non-porous surfaces, add 4 mL/L of water. Treated surfaces must remain wet for 10 minutes. The activity of 3M Quat Disinfectant Cleaner has been evaluated in the presence of 5% serum and 400 ppm hard water by the AOAC Use-Dilution test and found to be effective against a broad spectrum of gram negative and gram positive organisms as represented by:

Pseudomonas aeruginosa	Salmonella schottmuelleri
Staphylococcus aureus	Streptococcus faecallis
Salmonella choleraesuis	Shigella dysenteriae
Escherichia coli	Brevibacterium ammoniagenes
Streptococcus pyogenes	Salmonella typhi
Klebsiella pneumoniae	Serratia marcescens
Enterobacter aerogenes	Pseudomonas cepacia

FUNGICIDAL PERFORMANCE: 3M Quat Disinfectant Cleaner is an effective fungicide against Trichophyton mentagrophytes (the athlete's foot fungus) when used on surfaces in areas such as locker rooms, dressing rooms, shower and bath areas, exercise facilities, at 4 mL/L.

MOLD AND MILDEW CONTROL: At 4 mL/L 3M Quat Disinfectant Cleaner will effectively inhibit the growth of mold and mildew and the odors caused by them when applied to hard, non-porous surfaces (as indicated in general instructions above). Allow to dry on surface and repeat when mildew growth returns.

VIRUCIDAL PERFORMANCE: At 4 mL/L use-level, 3M Quat Disinfectant Cleaner was evaluated in the presence of 5% serum and 400 ppm hard water for a contact time of 10 minutes at room temperature (20-25EC) against the following virus: Influenza, A/Brazil, Herpes Simplex Type 1 and Type 2 and Vaccinia found on inanimate, hard, non-porous environmental surfaces. Instructions for Cleaning and Decontaminating Against HIV-1 on Surfaces and Objects Soiled with Blood or Bodily Fluids: Personal Protection: When handling items soiled with blood or bodily fluids, wear disposable latex gloves, gowns, masks, and eye coverings. Cleaning Procedure: Thoroughly clean blood or bodily fluids from surfaces before apply the disinfect. Disposal of Infectious Materials: Blood, body fluids, cleaning materials and clothing should be autoclaved and disposed of according to provincial and local regulations for infectious waste disposal.

This product is to be diluted only with the 3M Twist'n Fill Chemical Dispenser.

PRECAUTIONARY STATEMENTS - KEEP OUT OF REACH OF CHILDREN. Danger -Corrosive to eyes do not get in eyes, on skin or on clothing. Harmful if swallowed. Avoid contamination of food. Container Disposal - Triple rinse (or equivalent) and disposal trash collection.

FIRST AID: In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse. If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution, if not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

TOXICOLOGICAL INFORMATION: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

NOTICE TO USER: This control product is to be used only in accordance with the directions on this label. It is an offense under the Pest Control Products Act to use a control product under unsafe conditions.

NOTICE TO BUYER: Seller's guarantee shall be limited to the terms set out on the label and, subject thereto. The buyer assumes the risk to persons or property arising form the use or handling of this product and accepts the product on that condition.

REFER TO MATERIAL SAFETY DATA SHEET FOR MORE INFORMATION

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))