

93.05.19

Plant Products

7.5% CAPTAN GREENHOUSE FUNGICIDE DUST

COMMERCIAL

GUARANTEE: Captan 7.5%

REG. NO.: 9724 Pest Control Products Act

5 kg

Read the Label Before Using

EXPOSURE TO CAPTAN MAY PRODUCE LONG-TERM HEALTH EFFECTS. TO
MINIMIZE EXPOSURE FOLLOW DIRECTIONS OUTLINED ON THIS LABEL.

NOTICE TO USER:

This control product is to be used only in accordance with the
directions on this label. It is an offence under the Pest Control
Products Act to use a control product under unsafe conditions.

Plant Products Co. Ltd.
BRAMALEA
ONTARIO, CANADA
L6T 1G1

DIRECTIONS FOR USE:

For Black Spot of Roses, Carnation Leaf Spot and Blight, Chrysanthemum Grey Mold Blight, and Septoria Leaf Spot: Apply at first signs of disease. Give even coverage of upper and lower leaf surfaces. Repeat at 7-10 day intervals as necessary. As a greenhouse preplant bench treatment for Damping-off and Root Rot diseases: Apply 1.2 kg per 100m². Work into 7 to 10 cm of soil before planting.

PRECAUTIONS:

Keep out of reach of children. Never handle material with bare hands. Use rubber gloves; do not use leather or cloth gloves. Avoid spilling on skin or clothing. Do not breathe dust. Wear a respirator and goggles while mixing, loading and spraying. When handling or applying, wear goggles and clothing that completely cover arms and legs to minimize exposure. Change contaminated clothing daily and wash thoroughly before re-use. Wash hands, face and arms thoroughly after handling captan and before drinking, eating or smoking. Shower after use or at the end of the work day. If an accident occurs, remove contaminated clothing and wash skin thoroughly with soap and water. Wash clothing and gloves before re-use. Always work out of the wind during mixing and loading operations. Keep unprotected persons out of operating area or vicinity where there may be drift.

FIRST AID INSTRUCTIONS:

In case of poisoning, call a physician immediately. If on Skin, remove contaminated clothing and wash skin thoroughly with soap and water. If in Eyes flush with water for 5-10 minutes and obtain medical attention. If swallowed, give the patient one to two glasses of water, and cause vomiting by giving one dose of syrup of ipecac. If patient does not vomit within 20 minutes, give a second dose. If syrup of ipecac is not available, give the patient one to two glasses of water and cause vomiting by inserting finger down the throat. Repeat with water until vomit fluid is clear. The patient should be lying down with head below the level of the feet. DO NOT TRY TO CAUSE VOMITING IF THE PATIENT IS UNCONSCIOUS OR IN A CONVULSIVE STATE.

DECONTAMINATION AND DISPOSAL:

Rinse the emptied container thoroughly and add the rinsings to the spray mixture in the tank. Follow provincial instructions

for any required additional cleaning of the container prior to its disposal. Make the empty container unsuitable for further use. Dispose of the container in accordance with provincial requirements. For information on the disposal of unused, unwanted product and the cleanup of spills contact the regional office of the Environmental Protection, Environment Canada.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))