

09-MAR-2000

WILSON

GRANULAR ANT & GRUB KILLER

KILLS SOIL INSECTS IN LAWNS, GARDENS AND  
ORNAMENTAL SHRUBS AND BUSHES.

ANTS - EARWIGS - CHINCH BUGS  
SOD WEBWORMS - SOW BUGS - CRANE FLIES

KEEP OUT OF REACH OF CHILDREN

READ THE LABEL BEFORE USING

DOMESTIC

WITH DURSBAN®

5 KG

GUARANTEE: Chlorpyrifos 0.5%  
(DURSBAN(R))

REG. NO. 20792 P.C.P. ACT

Nu-Gro Ip Inc.  
10 Craig St.  
Brantford, Ontario  
Canada N3R 7J1

[www.wilson.nu-gro.com](http://www.wilson.nu-gro.com)

#### DIRECTIONS FOR USE

##### TURF AND LAWNS

Apply with a granular fertilizer spreader for uniform distribution. Apply one-half of required amount of product in one direction and the balance at right angles. Wash spreader thoroughly after use. Make no more than 2 applications per season.

White grubs, European chafer larvae, Japanese beetle larvae. Apply 400 g per 10 m<sup>2</sup> and water into turf. Apply when grubs are small and feeding near soil surface, usually in late July or August. Water in thoroughly.

Ants-Apply 200 g/10 m<sup>2</sup>. Treat ant nests and trails. Chinch

Bugs-Apply 200 g/10 m2 at first sign of damage. Water thoroughly. Keep off lawn until dry.

Sod Webworms-Apply 200 g/10 m2 at first sign of damage. Delay mowing for 12 to 24 hours after treatment.

Crane Fly Larvae (Leather Jackets) - Apply 200 g/10 m2 in late fall after the flight of adult crane flies has ceased for the year. Water in well.

#### ORNAMENTAL SHRUBS AND BUSHES

Ants, crickets, earwigs, sowbugs, spiders, cutworms, white grubs, European chafer larvae and Japanese beetle larvae. Apply to turf or to soil around plants at a rate of 400 g/10 m2. Make no more than 2 applications per season.

#### GARDENS

For control of cutworms and root maggots apply if damage is expected. Use on broccoli, Brussels sprouts, cabbage, Chinese cabbage, cauliflower, carrots, celery, cucumber, green pepper, onion, potato, sweet corn and tomato. Apply 200 g/10 m2 of soil surface before planting. Do not make more than one application per year. Rake soil lightly to incorporate granules into top 5 cm of soil. Do not use on green bunching onions.

Days to wait between application and harvest:

#### Crop

Broccoli.....	32
Brussels Sprouts.....	32
Cabbage.....	32
Carrot.....	60
Cauliflower.....	32
Celery.....	70
Chinese Cabbage.....	32
Cucumber.....	60
Green Pepper.....	40
Onion.....	50
Potato.....	7
Sweet Corn.....	70
Tomato.....	70

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN. Do not store granules with food or feed or drinking water. Do not use in poultry houses or other animal buildings. May cause irritation to skin or eyes; avoid contact. Avoid inhaling dust from granules. Wash thoroughly with soap and water after using.

WARNING: This product is toxic to fish and crustaceans. Avoid contamination of ornamental fish ponds and fish bowls.

FIRST AID: If swallowed, induce vomiting by giving patient 1 or 2 glasses of water and inserting finger down throat. Do not give anything by mouth to an unconscious person. Obtain medical attention or call a poison control centre. Flush contaminated eyes and skin with plenty of water for 15 minutes. For eyes get prompt medical attention.

TOXICOLOGICAL INFORMATION: Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically. Atropine by injection only is antidote.

DISPOSAL: Dispose of empty container in household garbage. Do not reuse.

®Dursban is a registered trademark of Dow AgroSciences LLC.

For additional product information or in case of emergency, spills or fire call toll-free 1-800-268-2806.

\*\*\*\*\*

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))