

90.07.19

((LABEL))

POAST C.T.

TANK MIX LIQUID HERBICIDE

Tank Mix Liquid Herbicide concentrate for postemergence control of annual grasses, wild oats and Canada thistle (top growth control) in canola (rapeseed).

Commercial (Agricultural)

GUARANTEE: Sethoxydim - 184 g/L
Clopyralid - 200 g/L

SPLIT CONTAINER

REGISTRATION NO.: 21015 PEST CONTROL PRODUCTS ACT

READ ENTIRE LABEL (INCLUDING ATTACHED PAMPHLET) BEFORE USING.

CAUTION POISON

DANGER: EYE IRRITANT

Net Contents:
8.6 L

BASF Canada Inc.
345 Carlingview Drive
Toronto, Ontario
M9W 6N9

BASF Canada Inc.
Suite 700
333-5th Avenue S.W.
Calgary, Alberta
T2P 3B6

Poast(R) and Assist(R) are registered trademarks of BASF AG.
BASF Canada Inc. is the registered user.

Caution: Read attached brochure for detailed application instructions before using.

PRECAUTIONS: 1. Keep out of reach of children. 2. Do not take internally. 3. Danger - eye irritant. May cause skin irritation. Avoid inhalation of vapour, dust or spray mist and contact with eyes, skin or clothing. 4. Wash thoroughly after handling and before eating, drinking or smoking. 5. Wear protective equipment and clothing, including goggles or face shield, gloves (rubber, PVC, neoprene or nitrile), hat, long-sleeved shirt, trousers and rubber boots. 6. If clothing becomes contaminated, remove and wash separate from household laundry before reuse. 7. Do not contaminate domestic or irrigation water, lakes, streams or ponds by the cleaning of equipment or the disposal of wastes. 8. Do not apply where run-off or erosion is likely to occur.

FIRST AID INSTRUCTIONS: IN CASE OF EYE CONTACT: Flush immediately with clear water for at least 15 minutes. Consult a physician if irritation develops. IN CASE OF SKIN CONTACT: Wash immediately with soap and water. Remove contaminated clothing and launder before reuse. Consult a physician if irritation develops. IF SWALLOWED: Give water or milk. Do not induce vomiting. Consult a physician immediately. IF INHALED: Remove person to fresh air. Assist breathing if necessary. Consult a physician immediately. Note to Physician: No specific antidote. Supportive care. Treatment based on judgement of the physician in response to reactions of the patient.

STORAGE: 1. Store in original, tightly-closed container. 2. Do not ship or store near food, feed, seed, fertilizers or pesticides. 3. Store in a cool, dry, locked, well-ventilated area without floor drain. 4. Herbicides should be shipped or stored separate from other pesticides to avoid cross-contamination. 5. Store in heated storage. Should product freeze, warm to room temperature and shake well before using.

DISPOSAL: 1. Triple-rinse the emptied container thoroughly and add the rinsings to the spray mixture in the tank. 2. Follow provincial instructions for any required additional cleaning of the container prior to its disposal. 3. Make the empty container unsuitable for further use. 4. Dispose of the container in accordance with provincial requirements. 5. For information on the disposal of unused, unwanted product and the clean up of spills, contact the regional office of Conservation and Protection, Environment Canada.

NOTICE TO USER: This control product is to be used only in accordance with the directions on this label. It is an offence under the PEST CONTROL PRODUCTS ACT to use a control product under unsafe conditions.

NOTICE TO BUYER: Seller's guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

((PAMPHLET))

POAST C.T.

TANK MIX LIQUID HERBICIDE

Tank Mix Liquid Herbicide concentrate for postemergence control of annual grasses, wild oats and Canada thistle (top growth control) in canola (rapeseed).

Commercial (Agricultural)

GUARANTEE:

Sethoxydim - 184 g/L

Clopyralid - 200 g/L

SPLIT CONTAINER

REGISTRATION NO.: 21015

PEST CONTROL PRODUCTS ACT

READ ENTIRE LABEL BEFORE USING.

DO NOT SHIP OR STORE WITH FOOD, FEEDS, DRUGS OR CLOTHING.

EMPTY THE ENTIRE CONTENTS OF THIS CONTAINER.

CAUTION

POISON

DANGER: EYE IRRITANT

BASF Canada Inc.
345 Carlingview Drive

Toronto, Ontario
M9W 6N9

BASF Canada Inc.
Suite 700
333-5th Avenue S.W.
Calgary, Alberta
T2P 3B6

GENERAL INFORMATION:

Poast C.T. tank mix concentrate is a broad- spectrum liquid herbicide designed to provide enhanced convenience for the control of annual grasses, wild oats and Canada thistle (top growth control) in canola (rapeseed).

The contents of this container will treat 4.0 hectares.

Empty the entire contents of this container.

TIMING OF APPLICATION:

WILD OATS AND ANNUAL GRASSES:

Apply when wild oats, green and yellow foxtails (wild millet), and other annual grasses, including barnyard grass, large crabgrass, volunteer corn, proso millet, fall panicum, witchgrass and Persian darnel, are in the 1 to 6 'true leaf' stage. Most effective control is achieved when application is made at the 2 to 5 'true leaf' stage when annual grasses are small (ie., plant height 10-15 cm) and actively growing, soil moisture is good, and the crop is small enough to permit thorough spray coverage. The preharvest interval is 70 days.

CANADA THISTLE:

For low to medium infestations (less than 10 plants per square metre) of Canada thistle, this product will provide top growth control for 6-8 weeks. Some regrowth will occur by the end of the season but this will not interfere with the harvesting of the crop.

Apply when Canada thistle is in the rosette to pre-bud stage of growth and at the 3-6 leaf stage of canola (rapeseed). Best results are obtained when Canada thistle is actively growing and soil moisture is adequate for rapid growth. Under dry soil

conditions, control of Canada thistle may be severely reduced. Applications made after the flowering stage of Canada thistle will not provide satisfactory control.

ADDITIVES AND WATER VOLUME:

Add Assist oil concentrate at a rate of 1.0 L/ha with the contents of this container to the spray tank in the manner outlined as follows.

MIXING:

1. Thoroughly clean the sprayer by flushing the system with water containing detergent. Trace contamination of the spray solution with 2,4-D, MCPA or similar herbicides will result in damage to the canola.
2. Fill clean spray tank with one half of the intended water to be used for spraying and start agitation system.
3. Empty the entire contents of this container into the spray tank to ensure uniform tank mixing.
4. Add sufficient quantity of Assist oil concentrate to obtain an application rate of 1.0 L/ha.
5. Add the remaining quantity of water necessary to obtain the desired total volume.
6. Continue to agitate or run the by-pass system.
7. After any break in the spraying operation, agitate thoroughly before spraying again. Check inside the tank to ensure that sprayer agitation is sufficient to re-mix the spray materials. Do not allow the mixture to sit overnight. Settling will occur if agitation is not continuous.
8. If an oil film starts to build up in the tank, drain it and clean the tank with a strong detergent solution.
9. Immediately after use, thoroughly clean the sprayer by flushing the system with clean water containing detergent.

NOTE: Failure to comply with these mixing instructions may result in poor weed control or crop injury.

SPRAYING INSTRUCTIONS:

Use sprayers equipped with standard flat fan nozzles. The use of 80' stainless steel flat fan nozzles is recommended for optimum spray coverage with nozzles tilted 45' forward to ensure better coverage.

The use of flood jet or hollow cone nozzles is not recommended because of uneven and inadequate spray coverage. Ensure thorough and uniform spray coverage over the entire leaf area of the target weeds.

Thoroughly clean all screens to prevent nozzle clogging.

Use a water volume of 100-200 litres per hectare at a pressure of 275 kPa.

RESTRICTIONS AND LIMITATIONS:

- . Do not apply by air.
- . Use only on canola (rapeseed).
- . When grasses are stressed due to drought, flooding or prolonged hot or cool temperatures (15'C or less), control can be reduced or delayed since grasses are not actively growing. Grass escapes or re-tillering may occur under prolonged stress conditions or low fertility.
- . Do not make applications to grasses stressed longer than 20 days due to lack of moisture as unsatisfactory control can result.
- . Thorough pre-plant tillage operations are required to fields where sod or forage grass crops may have grown in the previous year.
- . Rainfall within one hour of application may reduce the effectiveness of the spray.
- . Do not mix or apply this mixture with any other additive, pesticide or with fertilizer, except as specifically recommended on this label.
- . Allow 4 days between application of Poast C.T. and any other chemical.

- . Allow a minimum of 14 days between application of Poast C.T. plus Assist oil concentrate and re-planting of cereal or grass crop. A cultivation to a minimum depth of 10 cm is recommended 7 days prior to seeding.
- . Do not graze or feed treated forage to livestock prior to crop maturity.

WARNING:

Injury to sunflowers, legumes, fruit or vegetable crops, flowers or other desirable broadleaf plants as well as grass cereal or corn crops and turf may occur if Poast C.T. is applied directly on or is allowed to drift onto these susceptible crops.

Do not apply when weather conditions may cause spray drift from treated fields to adjacent crops. Wash sprayer thoroughly after use to avoid damage to the next crop sprayed, especially if it is a cereal, forage or corn crop.

CROP ROTATION:

Fields previously treated with Poast C.T. brand herbicide can be seeded to wheat, oats, barley, rye, flax, canola or should be summerfallowed.

Do not seed to crops other than those listed above, in the year following treatment.

PRECAUTIONS:

1. Keep out of reach of children.
2. Do not take internally.
3. Danger - eye irritant. May cause skin irritation. Avoid inhalation of vapour, dust or spray mist and contact with eyes, skin or clothing.
4. Wash thoroughly after handling and before eating, drinking or smoking.
5. Wear protective equipment and clothing, including goggles or face shield, gloves (rubber, PVC, neoprene or nitrile), hat, long-sleeved shirt, trousers and rubber boots.
6. If clothing becomes contaminated, remove and wash separate

from household laundry before reuse.

7. Do not contaminate domestic or irrigation water, lakes, streams or ponds by the cleaning of equipment or the disposal of wastes.
8. Do not apply where run-off or erosion is likely to occur.

FIRST AID INSTRUCTIONS:

IN CASE OF EYE CONTACT:

Flush immediately with clear water for at least 15 minutes. Consult a physician if irritation develops.

IN CASE OF SKIN CONTACT:

Wash immediately with soap and water. Remove contaminated clothing and launder before reuse. Consult a physician if irritation develops.

IF SWALLOWED:

Give water or milk. Do not induce vomiting. Consult a physician immediately.

IF INHALED:

Remove person to fresh air. Assist breathing if necessary. Consult a physician immediately.

Note to Physician:

No specific antidote. Supportive care. Treatment based on judgement of the physician in response to reactions of the patient.

STORAGE:

1. Store in original, tightly-closed container.
2. Do not ship or store near food, feed, seed, fertilizers or pesticides.
3. Store in a cool, dry, locked, well-ventilated area without floor drain.
4. Herbicides should be shipped or stored separate from other pesticides to avoid cross-contamination.
5. Store in heated storage. Should product freeze, warm to

room temperature and shake well before using.

DISPOSAL:

1. Triple-rinse the emptied container thoroughly and add the rinsings to the spray mixture in the tank.
2. Follow provincial instructions for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.
4. Dispose of the container in accordance with provincial requirements.
5. For information on the disposal of unused, unwanted product and the clean up of spills, contact the regional office of Conservation and Protection, Environment Canada.

NOTICE TO USER:

This control product is to be used only in accordance with the directions on this label. It is an offence under the PEST CONTROL PRODUCTS ACT to use a control product under unsafe conditions.

NOTICE TO BUYER:

Seller's guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

Poast(R) and Assist(R) are registered trademarks of BASF AG. BASF Canada Inc. is the registered user.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))