

91.08.13

DOMESTIC

250 mL TREATS 200 m2
1 Litre TREATS 800 m2

C-I-L
Lawn Weedkiller
TRICEP(R)

CAUTION POISON

KEEP OUT OF REACH OF CHILDREN

READ THE LABEL BEFORE USING

REG. NO. 10590 P.C.P. ACT

GUARANTEE: 2,4-D (present as amine) 100 g per litre
mecoprop d-isomer (present as amine) 50 g per litre
dicamba (present as amine) 9 g per litre

Chipman
A business Unit of C-I-L Inc.
Stoney Creek, Ontario
L8G 3Z1

C-I-L Lawn Weedkiller controls a broad spectrum of weeds including Dandelions, Plantain, Chickweed, Clover, Bedstraw, Buttercup, Black Medick (Yellow Clover), English Daisy, Healall, Creeping Charlie, Knotweed, Ragweed, Shepherd's Purse, etc.

DIRECTIONS:

5 mL = 1 teaspoon
15 mL = 1 tablespoon

Mix 125 mL with 10 L of water and spray evenly on 100 m2 of lawn. Apply in late spring or early fall when weeds are actively growing and temperatures are above 10°C. Do not cut grass for several days after application. Do not water within 24 hours. Avoid applying during a long, excessively dry or hot period or

when rain is expected. Do not apply to newly seeded lawns until the second or third mowing. Repeat as necessary. Fertilize the lawn to fill in bare spots after weeds are killed.

PRECAUTIONS:

Harmful if swallowed. Wash hands after use. Causes irritation of eyes and skin. Avoid inhaling spray mist and contact with eyes, skin or clothing. Avoid spray drift onto desirable plants. Do not use on newly seeded lawns or bent grass. Do not use clippings from a freshly sprayed lawn as a mulch. Do not use sprayer for other purposes. Do not store diluted spray. Avoid contamination of food, feed, fertilizer, seed, pesticides or any body of water. Store above 0°C. **Keep children and pets off treated area during application and until dry.**

DISPOSAL: Triple rinse container into sprayer and dispose of empty container with household garbage.

FIRST AID: If on skin, wash with soap and water. If in eyes, flush with water and get medical aid. If swallowed, obtain medical attention. In case of poisoning, call physician. Induce vomiting by giving patient one to two glasses of water and inserting a finger down the throat. Continue with water until vomit is clear. Patient should be lying down with head below level of feet. DO NOT TRY TO CAUSE VOMITING IF THE PATIENT IS UNCONSCIOUS OR IN A CONVULSIVE STATE.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))