

87.09.09

MAY & BAKER

NET CONTENTS 200 LITRES
METHOXOL 240-EC

INSECTICIDE-LIQUID EMULSIFIABLE CONCENTRATE

AGRICULTURAL

GUARANTEE:

METHOXYCHLOR 240 g/L

REGISTRATION NO. 14772

PEST CONTROL PRODUCTS ACT

READ THE LABEL BEFORE USING

WARNING POISON

MAY & BAKER CANADA INC.
6557 MISSISSAUGA ROAD
MISSISSAUGA, ONTARIO L5N 1A6

1274 PLAINS ROAD EAST
BURLINGTON, ONTARIO L7S 1W6

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN. Hazardous if swallowed, inhaled or absorbed through the skin. Avoid breathing vapours or spray mist. Use only in adequately ventilated area. Avoid contact with skin, eyes or clothing. After handling, wash thoroughly with soap and water before eating or smoking and change to clean clothing. Do not apply or allow to drift to areas occupied by unprotected persons and animals, or to streams, lakes or ponds to protect wildlife. Avoid contamination of feed or food, including such crops on which residue is unsafe.

FIRST AID: If swallowed, do NOT induce vomiting, but rush patient to nearest hospital taking this labelled container with you. In case of skin contact, remove contaminated clothing and wash skin thoroughly with soap and water. If in eyes, flush with plenty of water for 15 minutes and take patient to hospital at once.

TOXICOLOGICAL INFORMATION: Methoxychlor is an organochlorine

insecticide. Use saline cathartics and gastric lavages; avoid aspiration as this product contains petroleum distillates.

STORAGE: Do not store this product at temperatures below 20°C.

DISPOSAL:

1. Rinse the emptied container thoroughly and add the rinsings to the spray mixture in the tank.
2. Follow provincial instructions for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.
4. Dispose of the container in accordance with provincial requirements.
5. For information on the disposal of unused, unwanted product and the cleanup of spills contact the regional office of the Environmental Protection Service, Environment Canada.

NOTICE TO USER: This control product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products Act to use a control product under unsafe conditions.

NOTICE TO BUYER: Seller's guarantee shall be limited to the terms set out on the label and subject thereto the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

DIRECTIONS FOR USE

MIXING: Use oil as a diluent only if specifically recommended for a certain use. Oil will cause damage to trees and other ornamentals. The oil to be used in spray mixtures or solutions should be No. 2 Fuel Oil or No. 2 Diesel Oil. Use this table to prepare the spray mixture or solution of the required concentration.

12%	-	mix 1 part METHOXOL with	1 part of water
6%	-	mix 1 part METHOXOL with	3 parts of water
5%	-	mix 1 part METHOXOL with	4 parts of water or oil
2%	-	mix 1 part METHOXOL with	12 parts of water
1%	-	mix 1 part METHOXOL with	24 parts of water
0.5%	-	mix 1 part METHOXOL with	50 parts of water
0.25%	-	mix 1 part METHOXOL with	100 parts of water
0.1%	-	mix 2 part METHOXOL with	250 parts of water

HOST OR LOCATION AND INSECTS	METHOD OF APPLICATION	CONCENTRATION SEE MIXING TABLE
<p>ELM TREES Elm bark beetle (which transmits the Dutch elm disease) CAUTION: In cemeteries tombstones may be stained by spray drift.</p>	<p>MIST BLOWER APPLICATION: Apply 10 L per tree for a 15 elm (smaller trees in proportion) in late winter or early spring before budbreak.</p>	<p>12% spray - Mix with water only.</p>
	<p>A second application of 6% spray may be made in late July.</p>	<p>6% spray- Mix with water only</p>
	<p>HYDRAULIC SPRAYERS: Apply 100 L per tree when dormant.</p>	<p>2% spray - Mix with water only</p>
	<p>A second application of 1% may be made in late July.</p>	<p>1% spray-Mix with water only.</p>
<p>ORNAMENTAL TREES AND SHRUBS Caterpillars, Elm leaf beetle, Lace Bug, Leaf-hoppers, May beetle, Rose chafer, Tent caterpillar, Webworm, Weevils.</p>	<p>MISTBLOWER APPLICATION: Apply when insects are present. To be used by experienced pest control personnel only.</p>	<p>6% spray - Mix with water only.</p>
	<p>HYDRAULIC SPRAYERS: Apply on ornamentals when insects are present. Avoid contamination of fruit, vegetables and other edible crops.</p>	<p>0.1% spray - Mix with water only.</p>

<p>NON-CROP VEGETATION AND OTHER AREAS Adult mosquitoes, Mosquito larvae, Adult blackflies, Black fly larvae, Gnats</p>	<p>AIRCRAFT APPLICATION: Apply 4.25-5.5 L spray mixture per hectare (200-275 g actual methoxychlor). Use with care in farm ponds, fire ponds, stockponds or dugouts. Avoid treatment of streams, lakes or ponds where valuable fish may be harmed. For use in water wholly confined within the property of the user and where there is no outflow beyond the property limit.</p>	<p>5% spray -Mix with oil and water.</p>
<p>Adult mosquitoes, Adult blackflies, Adult blackflies, Gnats.</p>	<p>GROUND APPLICATION WITH THERMAL FOGGER: Apply Mix with oil 0.7-1.5 L of solution per hectare (35-70 g actual methoxychlor). Fogging does not give lasting residual control. Repeat this treatment as necessary.</p>	<p>5% solution -Mix with oil only.</p>
<p>BEEF CATTLE Hornflies.</p>	<p>BACKRUBBERS: Keep oilers moist for best results. Do not use within 7 days of slaughter. Do not use for dairy animals.</p>	<p>5% solution- Mix with oil only.</p>
<p>HORSES, SHEEP Stable flies, Mosquitoes, Fleas, Sheep ticks.</p>	<p>Spray horses at the rate of 30-60 mL per animal; repeat daily if necessary for control. Spray sheep at 30-60 mL per animal after shearing; repeat every two weeks. Do not apply within 7 days of slaughter. Use lower rate of spray on young horses and sheep. Never dip young animals less than 1 month old.</p>	<p>0.5%-1% spray- Mix with water only.</p>

MUSHROOM HOUSES Phorid and Sciarid Flies	BETWEEN CROPS: When house is empty spray inside walls, floors, windows, and woodwork to the point of run-off at the rate of 5L/100m ² . Repeat monthly until fall-filling begins. PRE-SPAWNING: Repeat application during cool-down when air temperature drops to 32°C. Do not spray ceilings or beds. PRE-CASING: Repeat 14 days after pre-spawning spray, no later than 4 days before casing. Do not spray ceilings or beds.	0.5% spray -Mix with water only.
	PRE-PRODUCTION: Repeat again 7 days after casing. Do not spray ceilings or beds but apply on side boards with a brush.	0.25% spray -Mix with water only.
COMPATIBILITY: METHOXOL 240-ED can be tank mixed with Cygon (R), Dimethoate or Malathion, and used as recommended on those labels.		

(R) Registered Trademark of Cyanamid Canada Ltd.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))