

22-JUL-1999

TomenAgro

COMMERCIAL

AGROX (R) I.F. Dual Purpose

Seed Treatment

Contains Diazinon and Captan

For the Control of Seed-Corn and Other Seed
and Root Maggots, Seed Rots, Seedling
Rots and Damping-Off Diseases.

WARNING

POISON

READ THE LABEL BEFORE USING

KEEP OUT OF REACH OF CHILDREN

CAUTION: Exposure to captan may produce long term health effects.
To minimize exposure, follow directions outlined on this label.

GUARANTEE:

diazinon 15%

captan 30%

REGISTRATION NO. 08349 PEST CONTROL PRODUCTS ACT

NET CONTENT: 10 kgs

TOMEN AGRO, Inc.
100 First Street, Suite 1700
San Francisco, CA 94105
415-536-3480
label number 20248-A

Distributor:
TOMEN CANADA, Inc.
Toronto Dominion Center, Suite 3312
Toronto, Ontario M5K 1K7
Canada
416-868-1103

AGROX I.F. Dual Purpose Seed Treatment is a specially formulated combination of diazinon for seed-corn maggot control, and captan for the control of fungus diseases of seedlings.

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN. Never handle material with bare hands. Use rubber gloves; do not use leather or cloth gloves. Avoid spilling on skin or clothing. Wash hands, face and arms thoroughly after handling this product and before drinking, eating or smoking. Do not breathe dust. When treating, augering or handling treated seed, work in a well ventilated area, and wear a suitable respirator, goggles and gloves. When handling or applying, wear goggles and clothing that completely covers arms and legs to minimize exposure. Change contaminated clothing daily and wash thoroughly before re-use. Shower after use or at the end of the work day. If an accident occurs, remove contaminated clothing, and wash skin thoroughly with soap and water. Wash clothing and gloves before re-use. Always work out of the wind during mixing and loading operations. Keep unprotected persons out of operating area or vicinity where there may be drift. Do not contaminate food or feed or any body of water. Label all unplanted treated seed: "Poisonous to man and animals. Do not use for food or feed. Do not sell to oil mills. This seed has been treated with captan and diazinon."

FIRST AID

In case of poisoning, call physician or a Poison Control Centre. If on skin, wash thoroughly with soap and water. if in eyes, flush with water for 15 minutes and get medical aid. If swallowed, give the patient one to two glasses of water and induce vomiting by inserting a finger down the throat. Continue with water until vomit is clear. Patient should be lying down with the head below the level of the feet. DO NOT TRY TO CAUSE VOMITING IF THE PATIENT IS UNCONSCIOUS OR IN A CONVULSIVE STATE.

TOXICOLOGICAL INFORMATION

Diazinon may cause cholinesterase inhibition. Atropine is antidotal.

EMERGENCY TELEPHONE NUMBER

All hours, 1-800-228-5635, ext. 174 ONLY for health and environmental information.

DISPOSAL

Thoroughly empty the contents of the container into the

application device. Make the empty container unsuitable for further use. Dispose of the container in accordance with provincial requirements. For information on the disposal of unused, unwanted product and the cleanup of spills, contact the provincial regulatory agency or the manufacturer.

DIRECTIONS FOR USE

Peas and Beans (all types including Lima, snap, field and soybeans) Corn and Sugar Beets - 52 g/25 g of seed. Apply AGROX I.F. Dual Purpose Seed Treatment as a dry application. For Slurry Machines - Mix 1.75 kg seed treatment per 10 L of water. If machine applies more or less slurry, adjust the amount of seed treatment so that amounts given above are applied to 25 kg of seed. For Hand Mixing - For every 25 kg of seed mix the given amount of seed treatment with 300 mL of water. Keeping mixture agitated and seed treatment in suspension in the water, pour over the seed and mix until uniformly coated. Allow to dry before seeding or bagging.

Potatoes - For the control of root maggots and disease, dip seed pieces in suspension of 325 g of AGROX I.F. Dual Purpose Seed Treatment per 100 L of water.

Onions and Radish - For the control of root maggots, apply 435 g of AGROX I.F. Dual Purpose Seed Treatment per kilogram of seed. Wet the seed with water containing 2% Methocel (methyl cellulose) and then add seed treatment. Mix thoroughly until well coated. Dry the seed before storing or planting.

NOTE: If the seed treatment increases the friction between seeds sufficient to reduce the rate of delivery through a grain drill, 30 g of graphite per 25 kg of seed should be added as a drill lubricant. Adjust the seed drill to give the required rate of seed delivery with the treated seed.

NOTICE TO BUYER

Seller's guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

NOTICE TO USER

This control product is to be used only in accordance with the directions on this label. It is an offense under the Pest Control Products Act to use a control product under unsafe conditions.

(R) Registered trademark of Tomen Agro, Inc.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))