

94.05.11

FLAMPROP-M

HERBICIDE

COMMERCIAL

Emulsifiable concentrate for control of Wild Oats

CAUTION POISON

WARNING IRRITANT

READ THE LABEL BEFORE USING

REGISTRATION NO. 23443
PEST CONTROL PRODUCTS ACT

GUARANTEE:

flamprop-m-methyl 52.5 g/L

NET: 20 L

UNITED AGRI PRODUCTS
R.R. #2
Dorchester, Ontario
N0L 1G5

DIRECTIONS FOR USE

GENERAL INFORMATION

FLAMPROP-M is a post-emergent herbicide for the control of wild oats in provincially recommended varieties of wheat (spring, durum, winter), triticale, canary grass, sunflowers and certain forage crops (seed production only).

TIMING OF APPLICATION

FLAMPROP-M can be applied to wild oats from the 3 leaf stage to the 6 leaf stage. For optimum control and maximum yield response, apply FLAMPROP-M when the majority of the wild oats

have 3-4 leaves.

INSERT ILLUSTRATION HERE

TIME AND RATE OF APPLICATIONS		
CROP	CROP STAGE	FLAMPROP-M Litres/Hectare
Wheat (provincially recommended varieties of spring, durum, winter), Canary grass, Triticale	Not beyond 6 leaf stage.	5
Sunflowers	Not beyond 6 leaf stage but before dense crop canopy forms	5 - 6.5
	Use the higher rate when wild oat pressure is heavy and for band application over crop row. Band application should be combined with interow cultivation.	
FORAGE CROPS FOR SEED PRODUCTION ONLY		
CROP	CROP STAGE	FLAMPROP-M Litres/Hectare
Alfalfa, Red clover, Birdsfoot Trefoil Sanfoin, Cicer milk vetch	In year of seeding	5
	Do not harvest or graze forages in year of treatment.	
Bromegrass, Meadow fescue, Creeping red fescue, Russian wild ryegrass, Intermediate wheatgrass, Crested wheatgrass	In year of seeding	5 - 7.5
	Use the high rate when grown without a companion crop. Use the low rate when grown with a companion crop for which FLAMPROP-M is registered. Do not harvest for forage or graze in year of treatment.	

MIXING INSTRUCTIONS

1. Fill the sprayer tank $\frac{1}{2}$ full with water and add the FLAMPROP-M. Engage gentle agitation.
2. Finish filling spray tank and maintain gentle agitation during mixing and spraying.
3. Do not let contents stand without agitation system engaged.

APPLICATION INSTRUCTIONS

Ground Application: Apply FLAMPROP-M in a minimum of 100 L/ha of water, using flat fan spray tips at a minimum pressure of 300 kPa. Use 50 mesh or larger in line and nozzle screens. Directing the spray forward 45° contributes to better coverage. If crop or weed canopy is dense, increase water volume to improve coverage and weed control.

Aerial application: Apply FLAMPROP-M by air with not less than 15 L/ha of water. Do not allow spray drift to contaminate other crops. Do not spray in winds greater than 15 km/hr. Application should not be made by air when oat crops are adjacent and downwind of the treated area.

PRECAUTIONS

Do not tank mix with any other herbicide. Do not tank mix with spray adjuvants, surfactants or liquid fertilizer. Allow 4 days between an MCPA or bromoxynil application and an application of FLAMPROP-M. Allow 7 days between a 2,4-D or dicamba application and an application of FLAMPROP-M. Rainfall within 2 hours of application may reduce effectiveness of FLAMPROP-M. Protect FLAMPROP-M from freezing.

KEEP OUT OF REACH OF CHILDREN. Product is harmful if swallowed. Avoid contact of this product with eyes. When handling, the use of safety goggles is recommended. Avoid inhalation of vapours. Keep product in original closed container and store away from food and feed. Wash hands and exposed skin before eating, drinking, smoking and after work. Flush concentrate from skin and eyes with water immediately after contact.

FIRST AID

This product contains petroleum distillate. INGESTED: DO NOT induce vomiting. If patient is unconscious, give him air. Keep patient prone and quiet. Transport the patient immediately to the nearest physician or contact a poison-

control centre. IF ON SKIN: Remove all contaminated clothing at once. Thoroughly wash skin with soap and water. Contact physician. Wash contaminated clothing before reuse. IF IN EYES: Flush immediately with running water for at least 15 minutes. Contact physician.

DISPOSAL

1. Rinse the emptied container thoroughly and add the rinsings to the spray mixture in the tank.
2. Follow provincial instructions for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.
4. Disposal of the container in accordance with provincial requirements.
5. For information on the disposal of unused, unwanted product and the clean-up of spills contact the regional office of Environmental Protection, Environment Canada.

NOTICE TO USER: This control product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products Act to use a control product under unsafe conditions.

NOTICE TO BUYER: Seller's guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))