



with plenty of water. Wash skin with warm water and soap. Get medical attention if irritation persists. Do not contaminate food or foodstuffs. Do not remain in treated areas. Vacate premises for at least 15 minutes after spraying. Ventilate the area after treatment is completed and before reentering. Remove pets and birds. Cover fish aquariums before spraying.

Food should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before use. When using the product in these areas, apply when the facility is not in operation.

Do not contaminate feed or water troughs. Any treated surface that may contact food or feed products should be washed thoroughly with potable water after spraying and before reuse. Do not use in areas where beekeeping equipment or supplies are stored.

Do not spray on plastic, painted or varnished surfaces or directly into electronic equipment such as radios, TVs, computers, etc.

FIRST AID: In case of contact with eyes or skin, flush with water. If irritation persists, seek medical attention. If accidentally swallowed, seek medical attention or call Poison Control Centre at once.

DISPOSAL: When container is empty, press valve to release all remaining pressure. Dispose of empty container in accordance with provincial requirements. For information on the disposal of unused, unwanted product and the cleanup of spills, contact the provincial regulatory agency or the manufacturer.

NOTICE TO USER: This Control Product is to be used only in accordance with the directions on the label. It is an offence under the Pest Control Products Act to use a control product under unsafe conditions.

NOTICE TO BUYER: Seller's guarantee shall be limited to the terms set out on the label and, subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

(( LEAFLET ))

CB (R)

PURGE (R)

CB-38

Insect Killer

COMMERCIAL

KILLS: House Flies, Barn Flies, Deer Flies, Stable Flies, Horn Flies, Mosquitoes, Gnats, Wasps, Hornets, Small Flying Moths, Cockroaches, Ants, Silverfish, Crickets, Centipedes, Sowbugs, Spiders, Indianmeal Moths, Angoumois Grain Moths, Confused Flour Beetles, Sawtoothed Grain Beetles, Red Flour Beetles, Rice Weevils, Granary Weevils, Grain Mites, Carpet Beetles, and Bedbugs.

READ THE LABEL BEFORE USING

GUARANTEE: Pyrethrins ..... 0.30%  
Piperonyl Butoxide, technical .... 2.40%

Product of U.S.A.

REGISTRATION NO. 23031  
PEST CONTROL PRODUCT ACT

CAUTION EXPLOSIVE

NET CONTENTS 652 g

Manufactured by:

WATERBURY COMPANIES, INC.  
P.O. Box 1812 - Waterbury, CT 06722 USA

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN. Do not use in presence of open flame or spark. Contents under pressure. Container may explode if heated. Do not expose to temperatures above 50°C. Do not puncture or incinerate.

Harmful if swallowed. Avoid breathing of vapor. Avoid contact with skin and eyes. In case of contact, immediately flush eyes with plenty of water. Wash skin with warm water and soap. Get medical attention if irritation persists. Do not contaminate food or foodstuffs. Do not remain in treated areas. Vacate premises for at least 15 minutes after spraying. Ventilate the area after treatment is completed and before reentering. Remove pets and birds. Cover fish aquariums before spraying.

Food should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before use. When using the product in these

areas, apply when the facility is not in operation.

Do not contaminate feed or water troughs. Any treated surface that may contact food or feed products should be washed thoroughly with potable water after spraying and before reuse. Do not use in areas where beekeeping equipment or supplies are stored.

Do not spray on plastic, painted or varnished surfaces or directly into electronic equipment such as radios, TVs, computers, etc.

FIRST AID: In case of contact with eyes or skin, flush with water. If irritation persists, seek medical attention. If accidentally swallowed, seek medical attention or call Poison Control Centre at once.

DISPOSAL: When container is empty, press valve to release all remaining pressure. Dispose of empty container in accordance with provincial requirements. For information on the disposal of unused, unwanted product and the cleanup of spills, contact the provincial regulatory agency or the manufacturer.

NOTICE TO USER: This Control Product is to be used only in accordance with the directions on the label. It is an offence under the Pest Control Products Act to use a control product under unsafe conditions.

NOTICE TO BUYER: Seller's guarantee shall be limited to the terms set out on the label and, subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

#### DIRECTIONS FOR USE

Hold container upright with nozzle away from you. Press valve down and spray as directed. Keep nozzle at least 90 cm from surfaces of fabrics to avoid possible staining. Remove pet, birds and cover fish aquariums before spraying.

Flies, Mosquitoes, Gnats, Wasps and Flying Moths: Close all doors and windows and spray room at the rate of 2 to 3 seconds per 30 m<sup>3</sup>. Direct spray to all parts of the room, especially windows and other light sources which attract these insects. Keep nozzle in constant motion. Keep room closed for 10 to 15 minutes after treatment. Do not remain in treated areas. Ventilate after treatment. Repeat treatment as necessary.

Cockroaches, Silverfish, Crickets, Spiders, Centipedes, and Sowbugs: Spray into hiding places such as cracks and crevices, behind sinks, cabinets, along baseboards and floors, around drains and plumbing, hitting insects with spray whenever

possible. Repeat treatment as necessary.

Ants: Spray trails, nests, and points of entry. Spray on ants when possible. Repeat as necessary.

Indianmeal Moths, Angoumois Grain Moths: Close doors and windows, direct spray into all parts of room, especially around stored food product containers, pallets, and darkened areas. Do not spray in areas where food is exposed in anyway. Spray 7 to 10 seconds for each 30 m<sup>3</sup> of space. Keep area closed for 15 minutes following treatment. Do not remain in treated areas. Ventilate after treatment. Repeat application weekly or as needed.

Exposed stages of Confused Flour Beetles, Sawtoothed Grain Beetles, Red Flour Beetles, Rice Weevils, Granary Weevils, Grain Mites: Spray into cracks and crevices, around cartons or containers and other areas where these insects tend to congregate. Spray 30 seconds for each 9.5 m<sup>2</sup> of surface. Keep area closed for 45 minutes to 1 hour after treatment. Repeat application as necessary.

Carpet Beetles: Spray directly on beetles, floors, baseboards and shelves. On carpets, use as a spot treatment only. Repeat treatment as necessary.

Bedbugs: Spray mattresses, particularly around tufts and seams. Take beds apart and spray into all joints. Treat baseboards, molding and floors. Repeat as necessary.

DAIRY FARM USE: Milking Parlor and Milk Room Close all windows and doors. Direct spray upwards, spraying for 5 seconds for 30 m<sup>3</sup> of space. Do not remain in treated areas. Ventilate after treatment. Remove milk and cover milking utensils before spraying. After spraying, milking utensils must be washed with an effective cleaning compound followed by a potable water rinse.

DAIRY CATTLE USE: Treat dairy cattle at least 20 minutes before milking, but do not spray directly on udder, wash udder before milking. Spray to slightly wet. Thoroughly spray entire animal from approximately 30 cm distance. Do not spray directly toward animal's eyes. Spot treat withers, shoulders and back where saliva accumulates from head tossing. Treat daily or as needed. Do not use more than once a day.

POULTRY OPERATIONS: In poultry houses, close all windows and doors. Spray at the rate of 5 seconds per 30 m<sup>3</sup> of space. Do not remain in treated areas. After 15 minutes, ventilate the area. Treat daily or as necessary.

HOG OPERATIONS: In hog houses, close all windows and doors. Spray at the rate of 2 to 3 seconds per 30 m<sup>3</sup> of space. Do not remain in treated area. After 15 minutes, ventilate the area. Repeat the application as necessary.

HOG USE: For animal application, direct spray over backs of hogs and spray to slightly wet. Do not spray directly toward the animal's face or eyes. Repeat application daily or as necessary. Do not use more than once a day.

STANCHION BARN USE: Walk behind animals and direct spray over backs and spray to slightly wet. For most effective results, apply each morning.

HORSE BARNS OR STABLES: In horse and pony barns, close all windows and doors. Spray at the rate of 5 seconds per 30 m3 of space. Do not remain in treated area. After 15 minutes, ventilate the area. Repeat the application as necessary.

HORSE USE: For animal application, lightly mist over backs of horses and ponies from a distance of 30 cm and spray to slightly wet. Do not spray directly toward the animal's face or eyes. Repeat application daily or as necessary. Do not use more than once a day. Do not cover horses after treatment.

BEEF CATTLE OPERATIONS: In barns, close all windows and doors. Spray at the rate of 5 seconds per 30 m3 of space. Do not remain in treated area. After 15 minutes, ventilate the area. Repeat application as necessary.

BEEF CATTLE USE: For animal application, thoroughly spray entire animal from approximately 30 cm distance and spray to slightly wet. Do not spray directly toward animal's face or eyes. Repeat application daily or as necessary.

FOOD PROCESSING PLANT USE: To kill Cockroaches, Spiders, Silverfish, Wasps, Hornets, Crickets Spray the insect directly when possible. Spray all probable hiding places such as cracks, crevices, baseboards, sinks, cabinets, shelves, under appliances, in food storage areas. To Kill Flies, Mosquitoes, Gnats, Small Flying Moths Close all windows and doors. Direct spray upward, spraying for 2 to 3 seconds for 30 m3 of space. Do not remain in treated areas. Ventilate after treatment.

\*\*\*\*\*

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))