

88.01.14

CLEAN CROP (R)

5% SEVIN DUST

CARBARYL INSECTICIDE

Controls many insects on  
VEGETABLES, ORNAMENTALS, LAWNS and POULTRY

COMMERCIAL

READ THE LABEL BEFORE USING

REGISTRATION NO. 18675 PEST CONTROL PRODUCTS ACT

GUARANTEE:

Carbaryl ..... 5%

YELLOWSTONE AGRI PRODUCTS  
12080 HORSESHOE WAY  
RICHMOND, B.C.  
V7A 4V5

NET CONTENTS: 10 kg

(R) CLEAN CROP is a Registered T.M. of United Agri Products Inc.

DIRECTIONS FOR USE: The low toxicity of SEVIN protects pets and animals and provides for safety in handling. SEVIN can be applied up to the day of harvest, unless otherwise indicated. Repeat at 7-10 day intervals as needed to maintain control.

VEGETABLES: The following amounts shown will cover 1 ha.

ASPARAGUS: To control Asparagus beetles, and climbing Cutworms, use 25-55 kg on the spears, and 60-110 kg on the ferns.

BEANS: To control climbing Cutworm, Lygus, and Stink bugs, use 50 kg; for Flea beetles, and Leafhoppers, use 40 kg and for Mexican bean beetles, use 28 kg.

BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER, AND KOHLRABI: To control Armyworms, Cabbage caterpillars, Cabbage loopers, Corn earworms, Lygus, and Stink bugs, use 37-55 kg; for Flea Beetles, and Leafhoppers, use 37-45 kg. Apply on a 7-10 day schedule when insect or damage appears. If cabbage loopers are the principle problem, use alternative insecticide after the heads begin to form.

EGGPLANT, PEPPER, POTATO, AND TOMATO: To control Colorado potato beetles, Leafhoppers, Potato flea beetles, Lygus and Stink bugs, Tomato fruitworms, and Tomato hornworms, use 25-45 kg.

PEAS: To control Alfalfa loopers, use 50 kg.

LEAFY VEGETABLES AND ROOT CROPS such as BEET, CARROT, CELERY, CHINESE CABBAGE, HORSERADISH, KALE, LETTUCE, PARSLEY, PARSNIP, RADISH, SPINACH, and TURNIP: To control Armyworms, Cabbage caterpillars, Cabbage loopers, Corn earworms, Lygus, and Stink bugs, use 37-55 kg; for Flea beetles, and Leafhoppers, use 37-45 kg.

LIMITATIONS: Do not apply within the following days before harvest.

- 1 day - asparagus, carrot, eggplant, pepper, and tomato;
- 3 days - bean, broccoli, brussels sprouts, cabbage, cauliflower, celery, cucumber, kohlrabi, head lettuce, pea, pumpkin, and squash;
- 7 days - beet (roots), horseradish, potato, radish, and turnip (roots);
- 14 days - beet (tops), Chinese cabbage, kale, leaf lettuce, parsley, parsnip, spinach, and turnip (tops).

ORNAMENTALS: To control Bagworms, Black vine weevils, Birch and Boxwood leaf miners, Boxelder bugs, Cankerworms, Gypsy moths, Japanese beetles, Lace bugs, Leaf aphids, Leaf beetles, Leafhoppers, Plant bugs, Psyllids, Rose aphids, Rose slugs, Scale insects, Tent caterpillars, and Thrips (exposed), apply a light, even coat to both sides of the foliage. Repeat as necessary. Some leaf injury may occur if applied when the foliage is tender or wet. Do not use on Boston ivy or Virginia creeper.

LAWNS and TURF: To control Ants, Chinch bugs, Climbing cutworms, Earwigs, Fall armyworm, Fleas, Leafhoppers, Millipedes, and Sod webworms (lawn moths), use 2-2.5 kg per 100 m<sup>2</sup>. Mow the lawn before application. Do not water for 2 days after treatment. Repeat in 2-3 weeks if necessary.

POULTRY such as CHICKENS, DUCKS, and TURKEYS: To control Lice and Northern fowl mites, and as a supplement to premise treatment for

Chicken mites, Fleas, and Fowl ticks, use 500 g per 100 birds. Dust, using the puff duster or shaker can. Direct the dust toward the neck and breast areas, and toward vent and fluff areas. Repeat in 4-8 weeks as required. Do not apply within 7 days of slaughter. Avoid contamination of eggs, feeds, nests, and water troughs. Do not treat birds having respiratory or other diseases which may be irritated by SEVIN DUST.

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN and Animals. Do not breathe the dust. Avoid skin contact. Wash hands before eating, smoking, or drinking. Avoid contamination of feed and food. Highly toxic to bees. Do not use where bees are foraging. Warn beekeepers to keep bees away from treated areas for one week after treatment.

FIRST AID: If swallowed, induce vomiting, by giving one dose (15 mL) of syrup of ipecac. If inhaled, move the patient to hospital.

TOXICOLOGICAL INFORMATION: CARBARYL is a moderate reversible cholinesterase inhibitor. Symptoms of poisoning: headache, nausea, vomiting, blurring of vision. A large dose of atropine may be needed to control the symptoms.

#### DISPOSAL

1. Follow provincial instructions for any required additional cleaning of the container prior to its disposal.
2. Make the empty container unsuitable for further use.
3. Dispose of the container in accordance with provincial requirements.
4. For information on the disposal of unused, unwanted product and the cleanup of spills, contact the regional office of Conservation and Protection, Environment Canada.

NOTICE TO USER: This control product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products Act to use a control product under unsafe conditions.

NOTICE TO BUYER: Seller's guarantee shall be limited to the terms set out on the label and, subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

\*\*\*\*\*

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label

information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))