

2000/2002

((leaflet))

HAGEN®

Flea Collar with Integral Buckle and
Choke Release Feature for Cats and Kittens

DOMESTIC

READ THE LABEL AND LEAFLET BEFORE USING.

DIRECTIONS: For best results, before placing this collar on, spray your pet with a pyrethrin based product such as Hagen Flea & Tick Spray for Cats. This will produce a "quick kill" effect.

Remove the collar from the package and place around cat's neck. Thread flat end of the collar through buckle hole and tighten as necessary, so that at least two ridges should be covered by the buckle. Ample room should be left to allow for growth. Cut off any excess length and discard in garbage. Collar must not be too tight or too loose (one finger should easily slip between collar and cat). The zig zag or perforated section of the collar allows stretching and release of the cat in many otherwise hazardous situations. Monitor cat or kitten and check fit frequently. After collar has been placed on cat, adult ticks will be killed in a few days and will fall off or may be easily removed. The collar will also control immature tick stages. Collar will kill ticks encountered in first 2 months of wear. Kills fleas for up to 4 months. FOR CONTINUOUS PROTECTION, REPLACE COLLAR WHEN EFFECTIVENESS DIMINISHES. To release the collar, squeeze between thumb and finger and pull sharply.

Temporary wetting will not reduce the effectiveness of this collar. For best results, use in combination with a complete flea eradication program (see your pet retailer for more information).

PRECAUTIONS: Do not open protective pouch until ready to use. DO NOT ALLOW CHILDREN TO HANDLE THIS COLLAR. Dust will form on the collar during storage. Do not get dust or collar in the mouth. Harmful if swallowed. Do not get dust in eyes. Wash hands thoroughly with soap and water after handling. Do not use on kittens younger than 12 weeks of age. When collar

is first worn, observe neck area frequently for signs of irritation. Remove collar immediately if irritation or adverse reaction is noted. Collar is intended for use only as an insecticide generator and is not to be taken internally by humans or animals. Do not use on sick or convalescing cats, or cats under drug treatment. Do not use other cholinesterase inhibitors (other organophosphate or carbamate containing parasiticides) on cat while collar is worn or for a few days after removing collar. Before using on pregnant animals, consult a veterinarian.

FIRST AID: If dust gets in eyes or on skin, flush with plenty of water. If irritation develops or persists, or if ingested, obtain medical attention.

TOXICOLOGICAL INFORMATION: Propoxur is a cholinesterase inhibitor. Atropine is antidotal.

DISPOSAL: Wrap and dispose of any excess collar, packaging and expired collars with garbage to keep out of reach of children.

Guarantee.....Propoxur 9.4%

REG. NO. 23543 P.C.P. ACT

NET WT: 12.5 g

DISTRIBUTED BY:
ROLF C. HAGEN INC.
3225 SARTELON
MONTREAL, QUEBEC H4R 1E8

Printed in Canada/Made in Germany

((box label))

HAGEN®

CONTENTS: 1 COLLAR/NET WT. 12.5 g(38.5cm)

Flea Collar with Integral Buckle
for Cats and Kittens

WITH CHOKE RELEASE FEATURE

3 STEP FLEA ATTACK PROGRAM
PREVENTATIVE TREATMENT

Kills Fleas for up to 4 months
And Brown Dog Ticks up to 2 months.

DOMESTIC - READ THE LABEL AND ENCLOSED LEAFLET BEFORE USING.

GUARANTEE: 9.4% PROPOXUR

REG. NO. 23543 P.C.P. ACT

Distributed by:
ROLF C. HAGEN INC.
3225 SARTELON
MONTREAL, QC H4R 1E8

Printed in Canada/Made in Germany

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))