

19-JUL-2000

((BOX))

DOMESTIC

250 mL

500 mL

C-I-L

Insecticide-Fungicide
for Roses and Flowers

FLORITECT (R)

READ THE LABEL AND BOX BEFORE USING

KEEP OUT OF REACH OF CHILDREN

GUARANTEE:

folpet	10.0%
carbaryl	10.0%
pirimicarb	2.5%

REGISTRATION NO. 12968 PEST CONTROL PRODUCTS ACT

ZENECA Agro
a business of ZENECA Corp.
Calgary, Alberta
T2E 7J2

Made in Canada

A99/G-4

KEEP THE BOX FOR USE DIRECTIONS

C-I-L Insecticide-Fungicide for Roses and Flowers has been specially formulated to control major diseases (black spot) and insects on roses, flowers and ornamentals.

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN. Harmful if swallowed. Wash thoroughly after use. Avoid inhaling spray mist or contact with eyes, skin or clothing. Do not apply indoors. Avoid contamination of food, feed, utensils, drinking water or any

body of water. Carbaryl is toxic to bees. Do not use where bees are foraging. Do not apply spray when temperature is over 27°C or under poor drying conditions. Keep children and pets away from treated area during application and until dry.

DISPOSAL

Triple rinse container into sprayer. Dispose of empty container and measuring cup with household garbage. Do not reuse empty container or measuring cup.

FIRST AID

If swallowed, induce vomiting by giving patient 1 or 2 glasses of water and inserting finger down throat. Call a poison control centre and obtain medical attention. If patient is unconscious, give him air. Do not give anything by mouth to an unconscious person. If in eyes, flush with water and obtain medical attention. If on skin, wash with soap and water.

TOXICOLOGICAL INFORMATION

Carbaryl and pirimicarb are moderate, reversible cholinesterase inhibitors. Atropine at 1-4 mg by intramuscular injections is antidotal followed by a further 2 mg every 30 min. as necessary. P2S and 2-PAM are not effective.

DIRECTIONS

5 mL = 1 teaspoon
15 mL = 1 tablespoon

WARNING: Some leaf injury may occur if treatments are made when tender foliage is wet or in the presence of prolonged high humidity. Do not use on Boston Ivy, Maidenhair Fern or Virginia Creeper.

SHAKE WELL BEFORE USING. Mix 10 mL per litre of water. Shake sprayer frequently while spraying. Thoroughly cover all sides of foliage and stems. Apply lightly on young tender foliage. Apply every 7 to 10 days or as needed.

Insects Controlled: Aphids, Caterpillars, Flea Beetles, Japanese Beetles, Lace Bugs, Leafhoppers, Leafrollers, Mites, Plant Bugs, Rose Aphids, Rose Chafer, Rose Slug, Sawflies, Scale insects (exposed crawler stage only), Thrips (exposed), Whiteflies.

Diseases Controlled: Black Spot of Roses, Powdery Mildew, Septoria Leaf Spot on Chrysanthemums, Anthracnose on Snapdragons, Alternaria Leaf Spot on Carnations, Marigolds and Zinnias.

Pirimicarb - Canadian Patent No. 933,159.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))