

03-MAY-2004 Notification received of label changes. See below for these changes.

07-FEB-2000

[Principal Display Panel]

FREEDOM™ WSB 75DF Herbicide

DRY FLOWABLE

For Use On Wheat (Spring, Winter or Durum), Barley, Oats,
Imazethapyr Tolerant Canola (e.g. Canola Varieties with the SMART
Trait), and CDC Triffid Flax

AGRICULTURAL

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING

GUARANTEE: Thifensulfuron methyl 75%

NET CONTENTS: 320 grams (4 x 80 gram water soluble bags)

REGISTRATION NO. 25935

PEST CONTROL PRODUCTS ACT

Do not sell water soluble bags separately.

~~DUPONT CANADA INC.~~
AGRICULTURAL PRODUCTS
BOX 2300, STREETSVILLE
MISSISSAUGA, ONTARIO L5M 2J4
1-800-667-3925

E.I. DU PONT CANADA COMPANY

Notification Change

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN.

Avoid breathing spray mist.

Avoid contact with skin, eyes and clothing.

Wear a long-sleeved shirt, long pants and chemical resistant gloves while handling this product.

Do not contaminate any body of water.

FIRST AID

EYE CONTACT: Flush eyes immediately with plenty of water for 15 minutes. Contact a physician or a poison control centre.

SKIN CONTACT: Wash skin with plenty of soap and water. Remove and wash contaminated clothing before reuse.

INGESTION: Administer water freely and induce vomiting. Repeat until vomit fluid is clear. Contact a physician or poison control centre immediately.

INHALATION: No specific intervention is indicated. Remove patient from area and contact a physician or poison control centre immediately.

For medical emergencies and transportation emergencies such as spills, leaks or fires, call 1-613-348-3616 (24 hours).

TOXICOLOGICAL INFORMATION

Treat Symptomatically

NOTICE TO USER

This control product is to be used only in accordance with the directions on this label. It is an offense under the PEST CONTROL PRODUCTS ACT to use a control product under unsafe conditions.

NOTICE TO BUYER

Seller's guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

The seller warrants that the purchase by the buyer and the use of this product, as such, will not infringe any Canadian patent.

STORAGE AND DISPOSAL

Storage:

Store product in original container only, away from other pesticides, fertilizer, food or feed. Not for use or storage in or around the home. Keep container closed. Keep water soluble bags in original bag away from moisture.

Disposal:

1. Make the empty container unsuitable for further use.
2. Dispose of the container in accordance with provincial requirements.
3. For information on the disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

[Directions for Use Brochure]

FREEDOM™ WSB 75DF Herbicide

DRY FLOWABLE

For Use On Wheat (Spring, Winter or Durum), Barley, Oats, Imazethapyr Tolerant Canola (eg. canola varieties with the SMART trait), and CDC Triffid Flax

AGRICULTURAL

READ THE LABEL AND THIS BOOKLET BEFORE USING

GUARANTEE: Thifensulfuron methyl 75%

REGISTRATION NO. 25935
PEST CONTROL PRODUCTS ACT

Do not sell water soluble bags separately.

~~DUPONT CANADA INC.~~
AGRICULTURAL PRODUCTS

E.I. DU PONT CANADA COMPANY

Notification Change

BOX 2300, STREETSVILLE
MISSISSAUGA, ONTARIO L5M 2J4
1-800-667-3925

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN.

Avoid breathing spray mist.

Avoid contact with skin, eyes and clothing.

Do not contaminate any body of water.

Wear a long-sleeved shirt, long pants and chemical resistant gloves while handling this product.

FIRST AID

EYE CONTACT: Flush eyes immediately with plenty of water for 15 minutes. Contact a physician or a poison control centre.

SKIN CONTACT: Wash skin with plenty of soap and water. Remove and wash contaminated clothing before reuse.

INGESTION: Administer water freely and induce vomiting. Repeat until vomit fluid is clear. Contact a physician or poison control centre immediately.

INHALATION: No specific intervention is indicated. Remove patient from area and contact a physician or poison control centre immediately.

For medical emergencies and transportation emergencies such as spills, leaks or fires, call 1-613-348-3616 (24 hours).

TOXICOLOGICAL INFORMATION

Treat Symptomatically

IMPORTANT

Injury to or loss of desirable trees or vegetation may result from failure to observe the following:

Do not apply, drain or flush equipment on or near desirable trees or other plants, or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots. Do not use on lawns, walks, driveways, tennis courts or similar areas. Prevent drift of spray to desirable

plants. Do not contaminate any body of water, including irrigation water that may be used on other crops.

Wheat, barley or oats must not be grazed by or fed to livestock within 7 days of application. Imazethapyr tolerant canola (e.g. canola varieties with the SMART trait) must not be harvested, grazed or fed to livestock within 64 days of application. CDC Triffid flax must not be harvested, grazed or fed to livestock within 82 days of application.

Carefully observe sprayer clean-up instructions, as spray tank residue may damage crops other than spring wheat, winter wheat, barley, oats, Imazethapyr tolerant canola (e.g. canola varieties with the SMART trait) and CDC Triffid flax.

DO NOT APPLY BY AIR.

GENERAL INFORMATION

FREEDOM™ WSB 75DF Herbicide is recommended for selective postemergence control of broadleaf weeds in wheat (spring, winter, or durum) barley, oats, Imazethapyr tolerant canola (e.g. canola varieties with the SMART trait) and CDC Triffid flax, not underseeded to legumes or grasses. FREEDOM™ WSB 75DF Herbicide is in the form of a dry flowable granule to be mixed in water and applied as a uniform broadcast spray. It is noncorrosive, nonflammable, nonvolatile, and does not freeze.

FREEDOM™ WSB 75DF Herbicide must be applied with a recommended surfactant early postemergence to the main flush of actively growing weeds. Warm, moist growing conditions promote active weed growth and enhance the activity of FREEDOM™ WSB 75DF Herbicide by allowing maximum foliar uptake and contact activity. Weeds hardened off by environmental stress may not be adequately controlled or suppressed and regrowth may occur.

FREEDOM™ WSB 75DF Herbicide may only be applied using ground equipment. For best results, ensure thorough spray coverage of target weeds. Refer to "Postemergence Applications" section of the label for complete use details.

FREEDOM™ WSB 75DF Herbicide stops growth of susceptible weeds rapidly. However, typical symptoms (discolouration) of dying weeds may not be noticeable for 1 to 3 weeks after application, depending on growing conditions and weed susceptibility. Favourable growing conditions following treatment promote the activity of FREEDOM™ WSB 75DF Herbicide while cold, dry conditions delay the activity. Degree of control and duration of effect depend on weed sensitivity, weed size, crop competition,

growing conditions and spray coverage.

DIRECTIONS FOR USE

POSTEMERGENCE APPLICATIONS

Wheat (spring, winter, or durum), barley or oats - not underseeded to legumes or grasses

Apply 20 grams of FREEDOM™ WSB 75DF Herbicide per hectare with a recommended surfactant such as AG-SURF® , AGRAL® 90, or CITOWETT® PLUS at 2 litres per 1000 litres spray solution (see "Mixing Instructions"), in the spring when the crop is in the two leaf to flag leaf (shotblade) stage of growth. Thorough coverage is important.

Weeds controlled:

Chickweed*
Hemp-nettle
Lady's-thumb
Redroot pigweed
Corn Spurry
Russian thistle
Green smartweed
Stinkweed
Lamb's-quarters
Volunteer rapeseed***
Cow cockle
Wild buckwheat**
Wild mustard

Weeds Suppressed:

Kochia

Suppression is a visual reduction in weed competition (reduced population or vigour) as compared to an untreated area.

For best results apply to emerged young actively growing weeds before the canopy closes. Weeds should be less than 10 cm tall or across. Weeds that emerge after treatment will not be controlled.

Note: If rain occurs soon after application control may be reduced. Several hours of dry weather are needed to allow FREEDOM™ WSB 75DF Herbicide to be absorbed by weed foliage.

One 80 gram water soluble bag of FREEDOM™ WSB 75DF Herbicide treats 4 ha.

Specific Weed Recommendations

* Chickweed: Apply FREEDOM™ WSB 75DF Herbicide plus recommended surfactant when the chickweed is small (1 to 6 leaf) and actively growing but before crop canopy prevents thorough coverage of weeds. Chickweed emerging after application will not be controlled.

** Wild buckwheat: Apply FREEDOM™ WSB 75DF Herbicide plus surfactant to actively growing wild buckwheat in the 1 - 3 leaf stage. Under dry conditions control may be reduced. Large plants may regrow after treatment.

*** Volunteer Rapeseed: FREEDOM™ WSB 75DF Herbicide will not control Imazethapyr tolerant canola (e.g. canola varieties with the SMART trait).

TANK MIXES

For use in Wheat (Spring, Winter or Durum), Barley, Oats

FREEDOM™ WSB 75DF HERBICIDE PLUS MCPA

FREEDOM™ WSB 75DF Herbicide may be tank mixed with MCPA Amine 500 to control the above weeds plus control of ragweed and suppression of Canada thistle.

Apply FREEDOM™ WSB 75DF Herbicide at 20 grams per hectare plus MCPA Amine 500 at 1.0 litre per hectare (or acid equivalent to deliver 500 grams active ingredient per hectare) plus a recommended surfactant such as AG-SURF® , AGRAL® 90 or CITOWETT® PLUS at 2 litres per 1000 litres of spray volume (see "Mixing Instructions" section of the label). Follow all precautionary statements for products used.

FREEDOM™ WSB 75DF Herbicide PLUS HOE-GRASS
HOE-GRASS 284 Herbicide is registered for use in Eastern Canada and British Columbia.

For use on Wheat (spring, winter and durum) and Barley (except Klages and Betzes varieties in Alberta).

Mix FREEDOM™ WSB 75DF Herbicide at 20 grams per hectare with HOE-GRASS 284 herbicide at 2.5 - 2.8 litres per hectare for the control of certain annual broadleaf weeds and wild oats only. Do not add a surfactant. This tank mix does not control green or yellow foxtail (wild millet, pigeon grass), Persian darnel, barnyard grass, or volunteer corn. Consult the HOE-GRASS® 284 label for additional application instructions and use precautions. Apply to broadleaf weeds that are young and

actively growing. Ensure that FREEDOM™ WSB 75DF Herbicide is completely in suspension in the spray tank before adding HOE-GRASS® 284. On repeat tank loads, prepare a FREEDOM™ WSB 75DF Herbicide/water slurry and add it to the required water volume in the spray tank prior to adding HOE-GRASS® 284.

FREEDOM™ WSB 75DF Herbicide PLUS AVENGE® 200-C

For control of broadleaf weeds and wild oats in certain varieties of wheat (check AVENGE® label for specific varieties) and all varieties of barley apply a tank mix of FREEDOM™ WSB 75DF Herbicide plus AVENGE® 200-C. Use 3.5 - 4.25 litres per hectare of AVENGE® 200-C with 20 grams FREEDOM™ WSB 75DF Herbicide per hectare. Consult the AVENGE® 200-C label for crop variety restrictions, additional application instructions, and precautions. DO NOT APPLY BY AIR. Applications of FREEDOM™ WSB 75DF Herbicide made at later stages of growth of broadleaf weeds may result in reduced control of these weeds. Ensure that FREEDOM™ WSB 75DF is completely in suspension in the spray tank before adding AVENGE® 200-C. Do not add a surfactant. On repeat tank loads, prepare a FREEDOM™ WSB 75DF Herbicide/water slurry and add it to the required water volume in the spray tank prior to adding AVENGE 200-C.

FREEDOM™ WSB 75DF Herbicide PLUS ACHIEVE 25 DG

For use on spring wheat and barley only.
FOR USE ONLY IN THE PRAIRIE PROVINCES AND THE PEACE RIVER REGION OF BRITISH COLUMBIA.

For control of broadleaf weeds and wild oats, and suppression of green foxtail in spring wheat and barley, apply 20 g/ha of FREEDOM™ WSB 75DF Herbicide tank mixed with 1 kg/ha of ACHIEVE® 25 DG. Add CHARGE at 10 litres per 1000 litres of spray solution (1% v/v). Apply when the crop is in the 2 leaf to flag leaf stage, and when the weeds are in the following stages:

Broadleaf weeds - less than 10 cm tall or across
Wild oats - 1-6 leaf (total leaves including tillers)
Green foxtail - 1-5 leaf (total leaves including tillers)

Consult the ACHIEVE® 25 DG label and directions for use for additional application instructions and use precautions. Do not apply within 60 days of harvest.

FOR USE ONLY ON:
IMAZETHAPYR TOLERANT CANOLA (E.G. CANOLA VARIETIES WITH THE SMART TRAIT) AND CDC TRIFFID FLAX

FOR USE ONLY IN THE PRAIRIE PROVINCES AND THE PEACE RIVER REGION

OF BRITISH COLUMBIA.

General Information

Imazethapyr Tolerant Canola (e.g. Canola varieties with the SMART™ trait)

Temporary crop yellowing may be observed shortly after application of the following treatments on Imazethapyr Tolerant Canola (e.g. Canola varieties with the SMART™ trait), especially when applied to the crop growing under environmentally stressful conditions (see USE PRECAUTIONS section of the label). Avoid application to Imazethapyr tolerant canola (e.g. canola varieties with the SMART™ trait) fields in which germination is uneven (some plants outside the recommended leaf stage for application), as crop injury may result. DO NOT apply to non-Imazethapyr tolerant canola as severe crop injury or death of the plant may occur.

CAUTION: Application of FREEDOM™ WSB 75DF herbicide on Imazethapyr tolerant canola (e.g. canola varieties with the SMART™ trait) may result in a delay in flowering.

FREEDOM™ WSB 75DF HERBICIDE ALONE

For Use on Imazethapyr Tolerant Canola (e.g. Canola Varieties with the SMART™ Trait) and CDC Triffid Flax

For the control of broadleaf weeds only, apply:

FREEDOM™ WSB 75DF herbicide at 20 grams per hectare plus a recommended surfactant such as AGRAL® 90, AG-SURF® or CITOWETT® PLUS at 2 litres per 1000 litres spray solution or the adjuvant SURE-MIX at 5 litres per 1000 litres spray solution (see "Mixing Instructions" section of label).

Apply FREEDOM™ WSB 75DF herbicide when the CDC Triffid flax is from 1 cm in crop height to the pre-bud stage.

Apply FREEDOM™ WSB 75DF herbicide when the Imazethapyr tolerant canola (e.g. canola varieties with the SMART™ trait) is in the 3 to 6 leaf stage of growth but prior to beginning of bolting. Application of FREEDOM™ WSB 75DF prior to the 3 leaf stage of Imazethapyr tolerant canola (e.g. canola containing the SMART™ trait) could result in crop injury.

FREEDOM™ WSB 75DF Herbicide will control the following broadleaf weeds:

WEEDS CONTROLLED

Chickweed*
Hemp-nettle
Lady's-thumb
Redroot pigweed
Corn spurry
Russian thistle
Green smartweed
Stinkweed
Lamb's-quarters
Volunteer rapeseed***
Cow cockle
Wild buckwheat**
Wild mustard

WEEDS SUPPRESSED

Kochia

Suppression is a visual reduction in weed competition (reduced population or vigour) as compared to an untreated area.

For best results apply to emerged young actively growing weeds before the canopy closes. Weeds should be less than 10 cm tall or across. Weeds that emerge after treatment will not be controlled.

Note: If rain occurs soon after application control may be reduced. Several hours of dry weather are needed to allow FREEDOM™ WSB 75DF to be absorbed by weed foliage.

Specific Weed Recommendations

* Chickweed: Apply FREEDOM™ WSB 75DF herbicide plus recommended surfactant when the chickweed is small (1 to 6 leaf) and actively growing but before crop canopy prevents thorough coverage of weeds. Chickweed emerging after application will not be controlled.

** Wild Buckwheat: Apply FREEDOM™ WSB 75DF plus surfactant to actively growing wild buckwheat in the 1 - 3 leaf stage. Under dry conditions control may be reduced. Large plants may regrow after treatment.

*** Volunteer Rapeseed: FREEDOM™ WSB 75DF will not control Imazethapyr tolerant canola (e.g. canola varieties with the SMART trait).

TANK MIXTURES IN:

IMAZETHAPYR TOLERANT CANOLA (EG. CANOLA VARIETIES WITH THE SMART

TRAIT) AND CDC TRIFFID FLAX.

FOR USE ONLY IN THE PRAIRIE PROVINCES AND THE PEACE RIVER REGION OF BRITISH COLUMBIA.

FREEDOM™ WSB 75DF + ASSURE® II

For Use on Imazethapyr Tolerant Canola (e.g. Canola Varieties with the SMART™ Trait) and CDC Triffid Flax

For the control of annual grasses as well as broadleaf weeds, apply a tank mixture of:

FREEDOM™ WSB 75DF	at 20 grams per hectare
plus	
ASSURE® II	at 0.38 or 0.5 litres per hectare
plus	
CANPLUS®™ 411	at 0.5-1.0 % v/v or 5-10 litres per 1000 litres spray solution
or	
SURE-MIX™	at 0.5 % v/v or 5 litres per 1000 litres spray solution

Apply the FREEDOM™ WSB 75DF plus ASSURE® II tank mixture when the CDC Triffid flax is from 1 cm in crop height to the pre-bud stage. Do not use on varieties of flax other than CDC Triffid.

Apply FREEDOM™ WSB 75DF plus ASSURE® II tank mixture when the Imazethapyr tolerant canola (e.g. canola varieties with the SMART trait) is in the 3 to 6 leaf stage of growth but prior to beginning of bolting.

Application of the FREEDOM™ WSB 75DF plus ASSURE® II tank mixture prior to the 3 leaf stage of Imazethapyr tolerant canola (e.g. canola varieties with the SMART™ trait) could result in crop injury.

This tank mix will control the above broadleaf weeds as well as the grassy weeds listed below:

RATE	WEEDS CONTROLLED	WEED LEAF STAGE
	Barnyard grass	2 leaf to early tillering *
	Green foxtail	2 leaf to early tillering

0.38 or 0.5

L/ha	Volunteer barley	2 leaf to early tillering
	Volunteer corn	2-6 leaf
	Volunteer oats	2 leaf to early tillering **
	Volunteer wheat	2 leaf to early tillering
	Wild Oats	1-5 leaf without tillers (0.38 L/ha)
		2 leaf to early tillering** (0.5 L/ha)

WEEDS SUPPRESSED

Quackgrass 2-6 leaf *

* Use the higher rate of ASSURE II (0.5 L/ha) for control of barnyard grass, as well as suppression of quackgrass.

** Best results on volunteer/wild oats if application is made before tillering begins.

Use SURE-MIX™ or the higher rate of CANPLUS® 411 when wild oats or quackgrass are the main target weeds, or under conditions not conducive to good growth.

NOTE: Consult the ASSURE® II label for additional instructions regarding grassy weed size and/or height at time of application as well as other use precautions and restrictions.

FREEDOM™ WSB 75DF + LONTREL™ 360 EC

For Use ONLY on Imazethapyr Tolerant Canola (e.g. Canola varieties with the SMART trait)

For control of annual broadleaf weeds plus top growth control of Canada Thistle, apply a tankmixture of the following:

FREEDOM™ WSB 75DF at 20 grams per hectare
 plus
 LONTREL™ 360 EC at 0.42 litres per hectare
 plus
 Non-ionic Surfactant at 0.2 % v/v
 or
 SURE-MIX™ at 0.5 % v/v or 5 litres per 1000 litres
 spray solution

Apply the FREEDOM™ WSB 75DF plus LONTREL™ 360 tank mixture when the Imazethapyr tolerant canola (e.g. canola varieties with the SMART trait) is in the 3 to 6 leaf stage of growth but prior to beginning of bolting. Application of the FREEDOM™ WSB 75DF plus LONTREL™ 360 tank mixture prior to the 3 leaf stage of Imazethapyr tolerant canola (e.g. canola varieties with the SMART

trait) could result in crop injury.

Weeds Controlled or Suppressed:

This tank mix will control or suppress the broadleaf weeds listed above with FREEDOM™ WSB 75DF used alone, PLUS provide top growth control of Canada Thistle. Consult the LONTREL™ 360 EC label for additional instructions regarding Canada Thistle size at the time of application, as well as other use precautions.

FREEDOM™ WSB 75DF + LONTREL™ 360 EC + ASSURE® II

For Use ONLY on Imazethapyr Tolerant Canola (e.g. Canola varieties with the SMART™ trait)

For top growth control of Canada Thistle as well as control of annual grasses and broadleaf weeds, apply a tank mixture of the following:

FREEDOM™ WSB 75DF at 20 grams per hectare
plus
LONTREL™ 360 EC at 0.42 litres per hectare
plus
ASSURE® II at 0.38 or 0.5 litres per hectare
plus
SURE-MIX™ at 0.5 % v/v or 5 litres per 1000 litres spray
solution
or
CANPLUS® 411 at 0.5 - 1.0 % v/v

Apply the FREEDOM™ WSB 75DF plus LONTREL™ 360 plus ASSURE® II tank mixture when the Imazethapyr tolerant canola (eg. canola varieties with the SMART™ trait) is in the 3 to 6 leaf stage of growth but prior to beginning of bolting. Application of the FREEDOM™ WSB 75DF plus LONTREL™ 360 plus ASSURE® II tank mixture prior to the 3 leaf stage of Imazethapyr tolerant canola (e.g. canola varieties with the SMART™ trait) could result in crop injury.

Weeds Controlled or Suppressed:

This tankmix will control or suppress the annual grasses & broadleaf weeds listed above with the FREEDOM™ WSB 75DF + ASSURE® II tankmix, PLUS provide top growth control of Canada Thistle. Consult the ASSURE® II and LONTREL™ 360 EC labels for additional instructions regarding weed size &/or height at the time of application, as well as other use precautions.

MIXING INSTRUCTIONS

1. Select a spray volume that will ensure thorough coverage and uniform spray pattern. Use a minimum of 100 L of spray volume per hectare.
2. Fill the spray tank 1/4 to 1/3 full with fresh water.
3. Add the recommended amount of FREEDOM™ WSB 75DF Herbicide to the spray tank while the agitator is running. Make sure all of the FREEDOM™ WSB 75DF Herbicide is in suspension. Continuous agitation is required to keep FREEDOM™ WSB 75DF Herbicide in suspension.
4. After FREEDOM™ WSB 75DF Herbicide is in suspension, other tankmix components may be added while continuing to fill the tank with water. The proper sequence for tank mix components is:

I. dry formulations

II. emulsifiable concentrates (i.e. MCPA, HOE-GRASS® , AVENGE® , LONTREL™ 360 EC or ASSURE® II)

III. surfactants or adjuvants

(recommended surfactants are AGRAL® 90, AG-SURF® , and CITOWETT® PLUS (recommended adjuvants for use with FREEDOM™ WSB 75DF +/- LONTREL™ 360 EC is SURE-MIX™ and with FREEDOM™ WSB 75DF + LONTREL™ 360 EC +/- ASSURE® II tankmixes are CANPLUS® and SURE-MIX™).

Note: Refer to the appropriate tank mix section of this label for more detailed recommendations on the specific surfactant or adjuvant required. A surfactant or adjuvant recommended on this label must be added to improve wetting and contact activity. Anti-foaming agents may be required.

Ensure that the herbicide is completely mixed before proceeding to the next step.

5. For repeat loads, reduce the tank heel to 10% or less of the previous load. Fill tank with fresh water as described in step 2, and continue as directed. Emulsifiable concentrates may make dispersion of FREEDOM™ WSB 75DF Herbicide more difficult. For that reason, tank heels of the previous tank mix should be kept to 10% or less of the spray tank volume.
6. Do not allow the spray mixture to remain in the tank for more than 24 hours before spraying or the effectiveness may be reduced. If the spray mixture has been allowed to stand in the tank, use vigorous agitation to thoroughly disperse before resuming spraying.

7. Use the spray preparation of FREEDOM™ WSB 75DF Herbicide within 24 hours or product degradation may occur. If spray solution is left standing without agitation, thoroughly agitate and re-suspend any settled FREEDOM™ WSB 75DF Herbicide before resuming the spraying operation.

NOTE: Do not mix the soluble bags with fertilizers, substances that contain boron or substances that release free chlorine. Mixing the water soluble bags with any of these compounds will result in an insoluble substance in the tank.

EQUIPMENT-SPRAY VOLUMES

Apply the spray mixture uniformly with properly calibrated ground equipment only. Ensure thorough coverage and a uniform spray pattern. Flat fan nozzles are recommended (minimum 100 litres spray volume per hectare). When using flood jet or "Raindrop" nozzles, apply 225 litres per hectare total spray volume to ensure thorough coverage. Use 50 mesh filter screens or larger (metal or nylon).

Continuous agitation is required to keep FREEDOM™ WSB 75DF Herbicide in suspension. Apply uniformly and avoid overlapping. Shut off spray boom while starting, turning, slowing or stopping to prevent crop injury from over application.

SPRAYER CLEANUP

To avoid subsequent injury to crops other than wheat, barley, oats, Imazethapyr tolerant canola (e.g. canola varieties with the SMART trait), or CDC Triffid flax immediately after spraying and prior to spraying other crops thoroughly remove all traces of FREEDOM™ WSB 75DF Herbicide from mixing and spray equipment as follows:

1. Drain tank and flush tank, boom and hoses with clean water for a minimum of ten minutes. Visually inspect tank to assure removal of all visible residues of FREEDOM™ WSB 75DF Herbicide. If necessary, repeat step 1. DO NOT CLEAN SPRAYER NEAR WELL OR WATER SOURCE OR NEAR TO DESIRABLE VEGETATION.
2. Fill tank with clean water while adding 1 litre household ammonia (containing a minimum 3% ammonia) per 100 litres of water. Flush solution through boom and hoses, and then add more water to completely fill tank. Allow to sit for 15 minutes with agitation. Again flush the hoses, boom and nozzles with the cleaning solution and drain tank.

3. Remove the nozzles and screens and clean separately in a bucket containing cleaning agent and water.
4. Repeat Step 2.
5. Thoroughly rinse the tank with clean water for a minimum of 5 minutes, flushing water through the hoses and boom.

CAUTION: Do not use ammonia with chlorine bleach. Using ammonia with chlorine bleach will release a gas with a musty chlorine odour which may cause eye, nose, throat, and lung irritation. Do not clean equipment in an enclosed area.

USE PRECAUTIONS

Do not apply to wheat, barley, oats, Imazethapyr tolerant canola (e.g. canola varieties with the SMART™ trait), or CDC Triffid flax that are or have been stressed prior to application by severe weather conditions (including frost or drought), low fertility, water saturated soil, disease or insect damage as crop injury may result. Do not apply to registered crops if the above stress conditions are expected within 3 days after application as crop injury may result.

Do not apply within 15 metres of a body of water or wetland area.

Do not apply to irrigated land where tail water will be used to irrigate crop land. Do not contaminate irrigation water.

Under certain environmentally stressful conditions (such as heavy rainfall, prolonged cool or hot weather, frost conditions, wide fluctuations in day/night temperatures), temporary lightening in crop colour and occasionally a slight reduction in crop height may occur.

Do not apply to crops underseeded to legumes, or grasses as injury to forage may result.

Because crop varieties differ in their tolerance to herbicides, limit first use of FREEDOM™ WSB 75DF Herbicide to a small area of each variety prior to adoption as a field practice.

Do not exceed a total of 20 g per hectare per crop year.

Do not plant any crop other than wheat, barley, Imazethapyr tolerant canola (e.g. canola varieties with the SMART™ trait) or CDC Triffid flax for 30 days after application of FREEDOM™ WSB 75DF Herbicide.

Some naturally occurring biotypes of certain weeds listed on this

label may not be controlled by ACCENT[®], ELIM[®], ULTIM[®], REFINE[®], ALLY[®], AMBER[®], EXPRESS PACK[®], MUSTER[®], PURSUIT[®], REFINE EXTRA[®], CLASSIC[®], PINNACLE[®], RELIANCE[™] STS , FREEDOM[®], TBM75, the CROSS[®] Toss-N-Go[®] component of CROSSFIRE[®], the EXTRA[®] component of CHAMPION EXTRA[®], the REFINE[®] component of REFINE[®] wo, the DF component of LASER[®] DF, or the PLUS component of TRIUMPH PLUS[®] and CHAMPION PLUS[®]. These populations are able to survive a use rate several times higher than needed to control susceptible populations. To delay selection for resistant populations rotate the use of ACCENT[®], ELIM[®], ULTIM[®], REFINE[®], ALLY[®], AMBER[®], EXPRESS PACK[®], MUSTER[®], PURSUIT[®], REFINE EXTRA[®], CLASSIC[®], PINNACLE[®], RELIANCE[™] STS[™], FREEDOM[™] , TBM75[®], CROSSFIRE[®], CHAMPION EXTRA[®], REFINE[®] wo, LASER[®] DF, TRIUMPH PLUS[®] and CHAMPION PLUS[®] with other herbicides effective on the same weeds. Avoid use of ACCENT[®], ELIM[®], ULTIM[®], REFINE[®], ALLY[®], AMBER[®], EXPRESS PACK[®], LASER[®] DF, MUSTER[®], PURSUIT[®], REFINE EXTRA[®], CLASSIC[®], PINNACLE[®], RELIANCE[™] STS[™], FREEDOM[™], TBM75, CROSSFIRE[®], CHAMPION EXTRA[®], REFINE[®] wo, TRIUMPH PLUS[®] and CHAMPION PLUS[®] repeatedly in the same field. Contact your local DuPont representative or the DuPont hotline at 1-800-667-3925 for further information.

NOTICE TO USER

This control product is to be used only in accordance with the directions on this label. It is an offense under the PEST CONTROL PRODUCTS ACT to use a control product under unsafe conditions.

NOTICE TO BUYER

Seller's guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

The seller warrants that the purchase by the buyer and the use of this product, as such, will not infringe any Canadian patent.

STORAGE AND DISPOSAL

Storage:

Store product in original container only, away from other pesticides, fertilizer, food or feed. Not for use or storage in or around the home. Keep container closed. Keep water soluble bags in original outer bag away from moisture.

Disposal:

1. Make the empty container unsuitable for further use.
2. Dispose of the container in accordance with provincial requirements.
3. For information on the disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

The DuPont Oval[®], CLASSIC[®], PINNACLE[®], STS[™], REFINE[®], REFINE EXTRA[®], ALLY , EXPRESS PACK[®], ASSURE[®], MUSTER[®], and SURE-MIX[™] are trademarks of E.I. du Pont de Nemours and Company. DuPont Canada Inc. is a licensee.

CHAMPION EXTRA[®], CHAMPION PLUS[®], CROSSFIRE[®], RELIANCE[™] and FREEDOM[™] are trademarks of DuPont Canada Inc.

ACHIEVE[®], CHARGE[®] and AGRAL[®] 90 are registered trademarks of the Zeneca Group of Companies.

AG-SURF[®] is a registered trademark of Interprovincial Cooperative Ltd.

AMBER[®] is a registered trademark of Novartis Crop Protection Canada Inc.

AVENGE[®], PURSUIT[®] and SMART[™] are trademarks of American Cyanamid Company.

CANPLUS[®] 411 is a registered trademark of ICI Americas Inc.

CITOWETT[®] PLUS is a registered trademark of BASF AG used under license by BASF Canada Inc.

HOE-GRASS[®] is a registered trademark of Hoechst Ag, Germany.

LASER[®] and TRIUMPH[®] Plus are registered trademarks of Hoechst Schering AgrEvo GmbH, Germany.

LONTREL[®] 360 is a trademark of Dow AgroSciences LLC.

[WATER SOLUBLE FILM LABEL]

FREEDOM WSB 75DF HERBICIDE

KEEP OUT OF REACH OF CHILDREN

CAUTION: SEE DIRECTIONS FOR USE FOLDER
FOR COMPLETE PRECAUTIONARY STATEMENTS

GUARANTEE: Thifensulfuron methyl 75%

DRY FLOWABLE

NET CONTENTS: 80 GRAMS

THIS BAG TREATS 4 HECTARES (10 ACRES)

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))