
GROUP	3	INSECTICIDE
-------	----------	-------------

**CYMBUSH® 250EC Pyrethroid Insecticide
Emulsifiable Concentrate**


AGRICULTURAL

GUARANTEE:

Cypermethrin250 g per litre

**READ THE LABEL AND ENTIRE LEAFLET BEFORE USING
KEEP OUT OF REACH OF CHILDREN**

WARNING



POISON

REGISTRATION NO. **19170**
PEST CONTROL PRODUCTS ACT

NET CONTENTS: **500 mL, 1 L, 5 L**

Syngenta Crop Protection Canada, Inc.
140 Research Lane
Research Park, University of Guelph
Guelph, Ontario
N1G 4Z3

H95/F-8

NOTICE TO USER

This control product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products Act to use a control product under unsafe conditions.

NOTICE TO BUYER

Seller's guarantee shall be limited to the terms set out on the label and, subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

CYMBUSH 250EC is a synthetic pyrethroid insecticide of low mammalian toxicity. It is a fast acting stomach and contact insecticide with no systemic or fumigant effect.

FIRST AID

IN CASE OF POISONING, contact a physician or a poison control centre **IMMEDIATELY**. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

DANGER - This product contains a petroleum distillate.

Therefore, **if swallowed**, do not induce vomiting. Get medical attention or call poison control centre immediately. If patient is unconscious, give air.

If in eyes, wash with plenty of clean water.

If on skin, wash with soap and water.

TOXICOLOGICAL INFORMATION

Treat symptomatically.

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN

- * Harmful or fatal if swallowed or absorbed through the skin.
- * Causes eye and skin irritation and skin sensitization. Avoid contact with skin, eyes and clothing.
- * Wear long-sleeved shirt or coveralls and gloves when handling or spraying. Wear face shield or eye goggles when mixing. Wash thoroughly after handling and before eating or smoking.
- * Avoid application when temperatures are above 27°C. Best control is obtained at cooler temperatures.
- * Cover crop or crop treated with CYMBUSH 250EC must not be used as a green feed for animals except as noted for corn, rutabagas and turnips.
- * **Do not apply by ground equipment within 15 m of productive fisheries water or waterfowl habitats.**
- * **For aerial application, an untreated border of 100 m must always be left around**

environmentally sensitive areas such as ponds, streams, rivers, dugouts and wetlands. If the 100 m buffer zone cannot be maintained, then ground application must be followed and the 15 m setback observed.

- * **Spray only when winds are less than 10 km per hour.**
- * Use spray nozzle tips and spraying pressure which provide small droplets.
- * Re-entry to treated areas is permissible as soon as spray deposit has dried.

If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in U.S., contact 1-866-375-4648 or www.cropro.org.

ENVIRONMENTAL HAZARDS

- * This product is very toxic to fish and aquatic organisms.
- * Do not apply when weather conditions favour drift from target area.
- * Do not contaminate ponds, lakes, streams or rivers during sprayer filling operation or while spraying.
- * This product is very toxic to bees; avoid spraying when bees are foraging. Spray deposit should be dry before bees commence foraging in treated crop.

This product contains a petroleum distillate which is moderately to highly toxic to aquatic organisms. Avoid contamination of aquatic systems during application. Do not contaminate these systems through direct application, disposal or waste or cleaning equipment.

STORAGE

Store in a cool dry well ventilated area away from foodstuffs and out of the reach of children and animals. This product is not affected by freezing.

SPILL CLEANUP

Wear appropriate protective equipment (gloves, glasses, apron) when attempting to clean up the spill. If the container is leaking, secure leak and place the container into a drum or heavy gauge plastic bag. Contact Syngenta Crop Protection Canada, Inc. (See EMERGENCY TELEPHONE NUMBER) for further information.

For spills and leaks; contain the liquid with dikes of inert material (soil, clay, kitty litter etc.). Absorb the spill onto inert material and shovel into a sealable waste container.

On hard surfaces - sprinkle spill area with detergent and scrub in a small quantity of water with a coarse broom. Let stand 10 minutes then absorb onto an inert material and shovel into the waste container.

On soil - remove the top 15 cm of soil in the spill area and replace with fresh soil. Dispose of all waste including scrub brush in accordance with provincial requirements.

For more information on the disposal of waste and the clean up of spills, contact the Provincial Regulatory Agency or the manufacturer. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

DECONTAMINATION AND DISPOSAL

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean up of spills.

CONTAINER DISPOSAL:

FOR DISPOSAL OF PLASTIC JUGS:

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

***IN CASE OF EMERGENCY INVOLVING A MAJOR SPILL, FIRE OR POISONING
CALL 1-800-327-8633 (FASTMED)***

GROUP **3** INSECTICIDE

CYMBUSH® 250EC Pyrethroid Insecticide Emulsifiable Concentrate

AGRICULTURAL

GUARANTEE:

Cypermethrin250 g per litre

**READ THE LABEL AND ENTIRE LEAFLET BEFORE USING
KEEP OUT OF REACH OF CHILDREN**

WARNING



POISON

REGISTRATION NO. **19170**
PEST CONTROL PRODUCTS ACT

NET CONTENTS: **500 mL, 1 L, 5 L**

Syngenta Crop Protection Canada, Inc.
140 Research Lane
Research Park, ~~University of Guelph~~
Guelph, Ontario
N1G 4Z3

Notification Change

NOTICE TO USER

This control product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products Act to use a control product under unsafe conditions.

NOTICE TO BUYER

Seller's guarantee shall be limited to the terms set out on the label and, subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

CYMBUSH 250EC is a synthetic pyrethroid insecticide of low mammalian toxicity. It is a fast acting stomach and contact insecticide with no systemic or fumigant effect.

FIRST AID

IN CASE OF POISONING, contact a physician or a poison control centre **IMMEDIATELY**. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

DANGER - This product contains a petroleum distillate.

Therefore, **if swallowed**, do not induce vomiting. Get medical attention or call poison control centre immediately. If patient is unconscious, give air.

If in eyes, wash with plenty of clean water.

If on skin, wash with soap and water.

TOXICOLOGICAL INFORMATION

Treat symptomatically.

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN

- * Harmful or fatal if swallowed or absorbed through the skin.
- * Causes eye and skin irritation and skin sensitization. Avoid contact with skin, eyes and clothing.
- * Wear long-sleeved shirt or coveralls and gloves when handling or spraying. Wear face shield or eye goggles when mixing. Wash thoroughly after handling and before eating or smoking.
- * Avoid application when temperatures are above 27°C. Best control is obtained at cooler temperatures.
- * Cover crop or crop treated with CYMBUSH 250EC must not be used as a green feed for animals except as noted for corn, rutabagas and turnips.
- * **Do not apply by ground equipment within 15 m of productive fisheries water or waterfowl habitats.**
- * **For aerial application, an untreated border of 100 m must always be left around**

environmentally sensitive areas such as ponds, streams, rivers, dugouts and wetlands. If the 100 m buffer zone cannot be maintained, then ground application must be followed and the 15 m setback observed.

- * **Spray only when winds are less than 10 km per hour.**
- * Use spray nozzle tips and spraying pressure which provide small droplets.
- * Re-entry to treated areas is permissible as soon as spray deposit has dried.

If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in U.S., contact 1-866-375-4648 or www.cropro.org.

ENVIRONMENTAL HAZARDS

- * This product is very toxic to fish and aquatic organisms.
- * Do not apply when weather conditions favour drift from target area.
- * Do not contaminate ponds, lakes, streams or rivers during sprayer filling operation or while spraying.
- * This product is very toxic to bees; avoid spraying when bees are foraging. Spray deposit should be dry before bees commence foraging in treated crop.

This product contains a petroleum distillate which is moderately to highly toxic to aquatic organisms. Avoid contamination of aquatic systems during application. Do not contaminate these systems through direct application, disposal or waste or cleaning equipment.

STORAGE

Store in a cool dry well ventilated area away from foodstuffs and out of the reach of children and animals. This product is not affected by freezing.

SPILL CLEANUP

Wear appropriate protective equipment (gloves, glasses, apron) when attempting to clean up the spill. If the container is leaking, secure leak and place the container into a drum or heavy gauge plastic bag. Contact Syngenta Crop Protection Canada, Inc. (See EMERGENCY TELEPHONE NUMBER) for further information.

For spills and leaks; contain the liquid with dikes of inert material (soil, clay, kitty litter etc.). Absorb the spill onto inert material and shovel into a sealable waste container.

On hard surfaces - sprinkle spill area with detergent and scrub in a small quantity of water with a coarse broom. Let stand 10 minutes then absorb onto an inert material and shovel into the waste container.

On soil - remove the top 15 cm of soil in the spill area and replace with fresh soil. Dispose of all waste including scrub brush in accordance with provincial requirements.

For more information on the disposal of waste and the clean up of spills, contact the Provincial Regulatory Agency or the manufacturer. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

DECONTAMINATION AND DISPOSAL

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean up of spills.

CONTAINER DISPOSAL:

FOR DISPOSAL OF PLASTIC JUGS:

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

***IN CASE OF EMERGENCY INVOLVING A MAJOR SPILL, FIRE OR POISONING
CALL 1-800-327-8633 (FASTMED)***

DIRECTIONS FOR USE

Ground Equipment Only (For Aerial Application See Page 7)

Precaution: Do not apply by ground equipment within 15 m of productive fisheries water or waterfowl habitats.

FRUITS (Apples, Pears, Peaches, Plums, Grapes, Strawberries): For all crops listed below, use sufficient water for thorough coverage. Follow provincial spray calendars for optimum timing of spray applications. Do not use more than 3 times per season. Leave a 15 metre buffer zone between water and sprayed area. **NOTE:** Repeated applications are not advised in orchards where integrated pest management is in effect because severe reductions of beneficial arthropods may result.

APPLES: Bud moth, Spotted Tentiform Leafminer, Leafrollers, Plum Curculio, Apple Maggot, Codling Moth - Apply at 400 mL/ha. For **Bud moth, Leafrollers, Plum Curculio and Codling Moth**, apply when insects or damage first appear. For **Spotted Tentiform Leafminer**, apply at peak emergence of the first and later generations of adults. Follow recommendations for the timing of sprays as given by local agricultural authorities based on insect monitoring. For **Apple Maggot**, apply first spray when adults appear in late June or early July. Repeat sprays every 10 days. CYMBUSH 250EC is not to be used more than 3 times during the season. Do not apply within 7 days of harvest. **Apple Leafcurling Midge and White Apple Leafhopper** - Apply at 400 mL/ha when insects or damage first appear. **Winter Moth** - Apply at 280 mL/ha when insects or damage first appear. CYMBUSH 250EC is not to be used more than 3 times during the season. Do not apply within 7 days of harvest.

Tank Mix with DIPEL - (Restricted for use in Nova Scotia only): Winter Moth, mix 28 mL of CYMBUSH 250EC with 560 mL of DIPEL and the recommended amount of spreader sticker in 600

to 800 L water and apply to one hectare. Apply once yearly at pre-bloom (half-inch green to pink stage) using an air blast orchard sprayer.

PEARS: Pear Psylla, Nymphs and Adults (Eastern Canada) and Codling Moth - Apply at 280 mL/ha. For **Pear Psylla**, follow spray calendar recommendations when pest first appears. **Pear Psylla (British Columbia) - Overwintering Adults:** Apply at 200 mL/ha when overwintering adult densities are highest. This usually occurs between silver tip and green tip. **Post Bloom Pear Psylla, Nymphs and Adults**, apply at 400 mL/ha when signs first appear or according to provincial spray calendar recommendations. CYMBUSH 250EC does not control the hard shell of Pear Psylla. Do not apply within 7 days of harvest.

GRAPES: Grape Leafhopper and Grape Berry Moth - Apply at 240 mL/ha. For **Grape Leafhopper**, apply when insects first appear. For **Grape Berry Moth**, apply shortly before bloom and repeat immediately after bloom. Additional sprays should be applied as insects or damage appear. CYMBUSH 250EC is not to be used more than 3 times per season. Do not apply within 7 days of harvest.

STRAWBERRIES: Strawberry Weevil, Tarnished Plant Bug and Meadow Spittlebug. For **Strawberry Weevil (Clipper)**, apply 280 mL/ha when buds are first visible and repeat when first buds show white. For **Tarnished Plant Bug**, apply 400 mL/ha at first bloom and repeat 7 to 10 days after first bloom. For **Meadow Spittlebug** apply 280 mL/ha when first buds show white. CYMBUSH 250EC is not to be used more than 3 times per season. See ENVIRONMENTAL HAZARDS for the protection of bees. Do not apply within 7 days of harvest.

PEACHES, NECTARINES: Oriental fruit moth, Tarnished plant bug and Oak Plant bug. Apply at 280 mL/ha with 550 L of water per hectare. Do not apply more than 2 applications. Do not apply within 7 days of harvest.

PLUMS: Plum Curculio - Apply at 400 mL/ha when insects or damage first appear. Do not apply more than 3 times per season. Do not apply within 7 days of harvest.

VEGETABLES

(Asparagus, Carrots, Celery, Cole Crops, Onions, Potatoes, Rutabagas, Tomatoes, Turnips): Apply at the rates indicated for each crop listed. Use sufficient water for thorough coverage unless the water volume is specified.

ASPARAGUS: Asparagus Beetle - Apply at 140 mL/ha to spears and ferns when insects are first observed. Repeat as required. Do not apply within 1 day of harvest.

CARROTS: Carrot Rust Fly - Apply at 280 mL/ha with 550 L of water per hectare when insects are first noticed. Follow recommendations for timing of sprays as given by local agricultural authorities based on insect monitoring. Do not apply more than 3 times per season. Do not apply within 35 days of harvest.

CELERY: Potato Leafhopper - Apply at 140 mL/ha in 500 L of water per hectare when insects are first noticed. Repeat as required up to 3 applications per season. Do not apply within 7 days of harvest.

COLE CROPS (Cabbage, Cauliflower, Broccoli, Brussels Sprouts): Cabbage Looper, Imported Cabbage Worm, Diamondback Moth Larvae - Apply at 140 mL/ha. Add AGRAL® 90 surfactant at 0.03% v/v (300 mL/1000 L of spray mix). Do not apply to muck soils. Use as needed at

2 week intervals up to a maximum of 3 applications per season. Leave a 3 day interval between the last spray and harvest. **Thrips** - Apply 200 mL/ha. Add AGRAL® 90 surfactant at 0.03% v/v (300 mL/1000 L of spray mix). Do not apply to muck soils. Use as needed at 2 week intervals up to a maximum of 3 applications per season. Leave a 3 day interval between the last spray and harvest.

CORN: (Sweet and Field): **European Corn Borer** - Apply at 280 mL/ha in 325 to 450 L of water per hectare using a boom and nozzle arrangement to ensure thorough spray coverage. Spray no later than when the first feeding is seen on foliage. Repeat sprays at 4 to 7 day intervals depending on the area and corn borer numbers (consult provincial agricultural representatives). Where there are two generations, late plantings of sweet corn will require sprays from the late whorl stage until close to harvest. Do not apply more than 3 times per season. Do not apply within 5 days of harvest. **Sweet corn plant refuse from processing plants and field corn silage derived from corn treated with CYMBUSH 250EC at the recommended rate and spray/harvest interval may be fed to lactating dairy cattle or beef cattle.**

LETTUCE: Aster Leafhopper - Apply 200 mL/ha beginning when damage or insects are first noticed. Apply up to 4 applications at 7-14 days interval. Do not apply within 14 days of harvest.

ONIONS: Onion Maggot Flies - To control onion maggots later in the growing season or on windrow onions prior to harvest, apply at 280 mL/ha. Follow recommendations for timing of sprays as given by local agricultural authorities based on insect monitoring. **Thrips** - Apply at 280 mL/hectare in sufficient water for good coverage. Apply at 10 day intervals as advised by local agricultural authorities. Do not apply more than 3 times per season. Do not apply within 3 days of harvest.

POTATOES: Colorado Potato Beetle, Potato Flea Beetle, Potato Leafhopper, Tuber Flea Beetle - Apply at 140 mL/ha when first signs of insects appear and damage is visible. Repeat as required up to a maximum of 3 applications per season. **Tarnished Plant Bugs:** Apply 200 mL/ha when insects appear. Repeat sprays as required. Do not apply within 7 days of harvest.

RUTABAGAS AND TURNIPS: Crucifer Flea Beetles - Apply at 200 mL/ha when insects appear and repeat as required. Do not apply within 21 days of harvest. Roots and tops may be fed to lactating dairy animals after the spray/harvest interval.

TOMATOES: Colorado Potato Beetle, Potato Flea Beetle, Potato Leafhopper - Apply at 140 mL/ha when first signs of insects appear and damage is visible. Repeat as required. Do not apply within 3 days of harvest.

FIELD CROPS:

Field Crops (Evening Primrose, Rapeseed, Mustard, Sunflowers and Tobacco) - Follow specific directions given for each crop listed. Unless otherwise indicated use sufficient water for thorough coverage.

EVENING PRIMROSE: Seed Weevil, Tarnished Plant Bug and *Microlepidoptera* spp. - Apply 280 mL in 300 L of water per ha at 2 to 3 week intervals when an insect population reaches an economic (significant) level. Apply a maximum of three applications per year.

RAPeseed (CANOLA) AND MUSTARD: Crucifer Flea Beetle - Apply 140 mL/ha when leaf injury is first noticed. **Bertha Armyworm** - Apply 200 to 280 mL/ha. Use higher rate for high infestations. Do not apply within 30 days of harvest.

SUNFLOWERS: Sunflower Beetle - Apply at 100 mL/ha in 100 to 120 L of water per hectare when insects first appear. Repeat if required. Do not apply within 70 days of harvest. **Sunflower Seed Weevil** - Apply at 100 mL/ha in 100 to 120 L of water per hectare when insects first appear. Repeat if required. Do not apply within 70 days of harvest.

TOBACCO: Cutworm (Dark-sided and White) Control: Tobacco (Greenhouse Plants): Mix 4.8 mL (1 teaspoonful) in 15 L of water and apply to 100 m² of plant bed. **Tobacco (Field):** For control of dark-sided cutworm and white cutworm, by methods given below, apply the recommended rate of CYMBUSH 250EC in 225 to 450 L of water per hectare using nozzle pressure of 175 to 350 kPa. Cutworm activity is greatest during the late evening and night. Application of CYMBUSH 250EC should be timed as close as possible to insect feeding activity. **Cover Crop Treatment:** Apply 140 mL per hectare once to rye or wheat cover when crop is 10 to 15 cm high, 4-5 days before ploughdown. Application should also be made to fence rows and to a 15 m strip into nearby cover crop. Allow 60 days between the last treatment and harvest in barley and 30 days on wheat. Do not use treated cover crops as green feed for animals. **Soil Treatment:** Apply 280 mL per hectare once to the soil 5 days before transplanting. Do not incorporate. Do not disturb the soil surface for at least 5 days following treatment since mixing of CYMBUSH 250EC with soil will reduce its effectiveness. Application should also be made to fence rows and to a 15 m strip into nearby cover crops. **Post Planting Treatment:** At transplanting, apply 280 mL per hectare of area sprayed in a 25 cm band over the row using 150 to 300 L of water per hectare. Under conditions of severe insect pressure, application should be made to fence rows and to a 15 m wide strip into nearby cover crops. A follow-up treatment may be necessary if there are late developing cutworms.

STEVIA: Cutworms-black, variegated and darksides - Apply at 280 mL/ha in sufficient water. Maximum of two applications per year. Ground application only. Apply post transplant as indicated by scouting. Do not apply within 50 days of harvest.

CONIFER SEEDLING (NURSERY): Tarnished Plant Bug (*Lygus sp*) - Apply 280 mL/ha in sufficient water for good coverage. Maximum of 3 applications per growing season. **NOT FOR GREENHOUSE USE.** Do not apply using back pack sprayers. Do not apply before stock thinning.

DIRECTIONS FOR USE

AERIAL APPLICATION

CYMBUSH 250EC may be applied once or twice per season as indicated for each crop listed below.

Specific Use Directions for Aerial Application

Read and understand the entire label before using this product.

Label rates, conditions and precautions are product specific. Apply only at the rate recommended for aerial application on this label. Where no rate for aerial application appears for a specific use, this product cannot be applied by any type of aerial equipment.

For aerial applications, ensure the aircraft is equipped and calibrated to deliver a uniform spray coverage with a minimum potential for drift. To ensure uniform application use an appropriate marking device. Apply in weather conditions that will not promote drift. Suggested conditions for good aerial application are moderate temperatures (less than 25°C), moderate relative humidity (greater than 40%), and light winds (3 to 9 kph). Aerial drift is increased under certain meteorological conditions. Do not apply during periods of dead calm, when winds are gusty or when temperature inversion is likely (e.g. evening when warm air is rising from the crop or morning when sunshine warms the soil and air rises from the field). Do not apply by air

in winds greater than 16 kph at flying height at the site of application.

Use only medium or coarse nozzles rated as delivering droplets of volume median diameter of 215 microns or greater. Examples: 6506 flat fan, CP 0.125 deflector nozzle with a low shear deflector angle (30 degrees), D12-56 disc-core.

Apply in a spray volume of 10 to 40 litres per hectare. Use water volumes at the higher end of this range to ensure good coverage for optimum insect control and to minimize drift. Do not apply more than recommended rates of CYMBUSH 250EC per hectare per season by air.

Do not apply to any body of water. Avoid drifting of spray onto any body of water or other non-target areas.

This product is very toxic to fish and aquatic organisms. For the protection of non-target habitats, overspray or drift to sensitive habitats must be avoided. A buffer zone of 100 metres is required between the downwind edge of the boom and the closest edge of sensitive aquatic habitats such as sloughs, ponds, coulees, prairie potholes, lakes, rivers, streams, and wetlands, (and wildlife habitat at the edge of these bodies of water). Do not contaminate these habitats when cleaning and rinsing spray equipment or containers.

Apply only under conditions of good practice specific to aerial application as outlined in the *Basic Knowledge Requirements for Pesticide Education in Canada: Applicator Core and Aerial Module*, developed by CAPCO.

Do not allow the pilot to mix chemicals to be loaded onto the aircraft. Loading of premixed chemicals with a closed system is permitted.

It is desirable that the pilot have communication capabilities at each treatment site at the time of application.

The field crew and the mixer/loaders must wear chemical resistant gloves, coveralls and goggles or face shield during mixing/loading, cleanup and repair. All personnel on the job site must wash hands and face thoroughly before eating and drinking. Protective clothing, aircraft cockpit and vehicle cabs must be decontaminated regularly.

Clean and decontaminate protective clothing and application equipment regularly.

Read and understand the entire label before opening this product. If you have questions, call the manufacturer or obtain technical advice from the distributor or your provincial agricultural representative. Application of this specific product must meet and/or conform to the following:

CORN (Sweet and Field): European Corn Borer - Apply at 280 mL/ha in 20 to 45 L of water per hectare. Spray no later than when the first feeding is seen on foliage. Repeat sprays at 4 to 7 day intervals depending on the area and corn borer numbers (consult provincial agricultural representatives). Where there are two generations, late plantings of sweet corn will require sprays from the late whorl stage until close to harvest. Up to two applications per season by air may be applied as needed. Do not apply within 5 days of harvest.

Sweet corn plant refuse from processing plants and field corn silage derived from corn treated with CYMBUSH 250EC at the recommended rate and spray/harvest interval may be fed to lactating dairy cattle or beef cattle.

POTATOES: Colorado Potato beetle, Potato Flea Beetle, Potato Leafhopper, Tuber Flea Beetle - Apply at 140 mL/ha in 20 to 45 L of water when first signs of insects appear and damage is visible. **Tarnished Plant Bug:** Apply 200 mL/ha in 20 to 45 L of water when insects appear. Repeat sprays as required. Up to two applications per season may be applied by air as needed. Do not apply within 7 days of harvest.

RAPSEED (CANOLA): Crucifer Flea Beetle - Apply 140 mL/ha when leaf injury is first noticed. **Bertha Armyworm** - Apply 200 to 280 mL/ha in 20 to 45 L of water. Use higher rate for high infestations. Only one application per season may be applied by air as needed. Do not apply within 30 days of harvest. Do not apply to mustard.

SUNFLOWERS: Sunflower beetle - Apply at 100 mL/ha in 20 to 45 L of water per hectare when insects first appear. Only one application per season may be applied by air as needed. Do not apply within 70 days of harvest.

Sunflower Seed Weevil - Apply at 100 mL/ha in 20 to 45 L of water per hectare when insects first appear. Only one application per season may be applied by air as needed. Do not apply within 70 days of harvest.

Resistance-Management Recommendations

For resistance management, please note that Cymbush 250EC contains a Group 3 insecticide. Any insect population may contain individuals naturally resistant to Cymbush 250EC and other Group 3 insecticides. The resistant individuals may dominate the insect population if this group of insecticides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action but are specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay insecticide resistance:

Where possible, rotate the use of Cymbush 250EC or other Group 3 insecticides with different groups that control the same pests in a field.

Avoid application of more than the maximum number listed in the label and consecutive sprays of Cymbush 250EC or other insecticides in the same group in a season.

Use tank mixtures with insecticides from a different group when such use is permitted.

Insecticide use should be based on an IPM program that includes scouting, record keeping, and considers cultural, biological and other chemical control practices.

Monitor treated pest populations for resistance development.

Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for the specific site and pest problems in your area.

For further information or to report suspected resistance, contact company representatives at 1-800-459-2422 (in Eastern Canada) or 1-800-665-9250 (in Western Canada) or at www.syngenta.ca.

05-FEB-1998

AGRICULTURAL1 L
5 L
500 mL

SUPPLEMENTAL LABEL

For the of black, variegated and
darksided cutworm on Stevia

CYMBUSH(R) 250EC

Pyrethroid Insecticide

Emulsifiable Concentrate

WARNING

POISON

READ THE LABEL AND ENTIRE LEAFLET BEFORE USING

KEEP OUT OF REACH OF CHILDREN

GUARANTEE: cypermethrin 250 g per litre

REGISTRATION NO. 19170 PEST CONTROL PRODUCTS ACT

ZENECA Agro, a business of ZENECA Corp.
250 - 3115 12th Street N.E.
Calgary, Alberta
T2E 7J2

CYMBUSH 250EC is a new synthetic pyrethroid insecticide of low mammalian toxicity. It is a fast acting stomach and contact insecticide with no systemic or fumigant effect.

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN

- * Harmful or fatal if swallowed or absorbed through the skin.
- * Causes eye and skin irritation and skin sensitization. Avoid contact with skin, eyes and clothing.
- * Wear long-sleeved shirt or coveralls and gloves when handling or spraying. Wear face shield or eye goggles when mixing. Wash thoroughly after handling and before eating or smoking.
- * Avoid application when temperatures are above 27°C. Best control is obtained at cooler temperatures.
- * Cover crop or crop treated with CYMBUSH must not be used as a green feed for animals except as noted for corn, rutabagas and turnips.
- * Do not apply by ground equipment within 15 m of productive fisheries water or waterfowl habitats.
- * For aerial application, an untreated border of 100 m must

always be left around environmentally sensitive areas such as ponds, streams, rivers, dugouts and wetlands. If the 100 m buffer zone cannot be maintained, then ground application must be followed and the 15 m setback observed.

- * Spray only when winds are less than 10 km per hour.
- * Use spray nozzle tips and spraying pressure which provide small droplets.
- * Re-entry to treated areas is permissible as soon as spray deposit has dried.

FIRST AID

DANGER - This product contains a petroleum distillate. Therefore, if swallowed, do not induce vomiting. Get medical attention or call Poison Control Centre immediately. If patient is unconscious, give air. If in eyes, wash with plenty of clean water. If on skin, wash with soap and water.

EMERGENCY TELEPHONE NUMBER

All hours, 1-800-263-0984 ONLY for health and environmental information.

DISPOSAL

Rinse the empty container thoroughly and add the rinsings to the spray mixture in the tank. Follow provincial instructions for any required additional cleaning of the container prior to its disposal. Make the empty container unsuitable for further use. Dispose of container in accordance with provincial requirements. For information on the disposal of unused, unwanted product and the cleanup of spills, contact the Provincial Regulatory Agency or the Manufacturer.

NOTICE TO USER

This control product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products Act to use a control product under unsafe conditions.

DIRECTIONS FOR USE

STEVIA: Cutworms-black, variegated and darksided - Apply at 280 mL/ha in sufficient water. Maximum of two application per year. Ground application only. Apply post transplant as indicated by scouting. Do not apply within 50 days of harvest.

NOTICE TO BUYER:

Seller's guarantee shall be limited to the terms set out on the label and subject thereto. The buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

CYMBUSH is a registered trademark of a Zeneca Group Company.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))