SANEX

LINDANE 10% E.C.

Emulsifiable Concentrate Insecticide for Control of Forest Pests, Pests on Animals and Pests Infesting buildings.

AGRICULTURAL

COMMERCIAL

WARNING

POISON

GUARANTEE: GAMMA BHC (LINDANE) 10%

READ THE LABEL BEFORE USING

NET CONTENTS: 500 mL

REGISTRATION NO. 6122 PEST CONTROL PRODUCTS ACT

Manufactured by:
SANEX INC.
5100 Timberlea Blvd.
Mississauga, Ontario
L4W 2S5

FIRST AID: In case of contact with eyes, flush eyes with plenty of water for at least 15 minutes and consult a physician immediately. For skin contact, wash with soap and water. Remove and wash contaminated clothing and shoes before re-use. Product contains petroleum distillate. If accidentally swallowed, do not induce vomiting. Seek medical attention at once.

PRECAUTIONS: WARNING: KEEP OUT OF REACH OF CHILDREN AND ANIMALS. Keep in original container during storage. Destroy empty container. Handle and apply only as recommended and at recommended rate. Do not contaminate streams, lakes, ponds, irrigation water, water used by livestock or water used for domestic purposes. Do not spray when plant is in operation. Do not spray on working surfaces or food handling equipment. Do not spray when food when food is exposed in the area. Do not use on calves or colts under 3 months of age. Do not use on sick, emaciated or dairy animals. Do not treat meat animals within 30 days before slaughter.

DISPOSAL: Rinse the emptied container thoroughly and add the rinsings to the spray mixture in the tank. Follow provincial instructions for any required additional cleaning of the container prior to its disposal. Make the empty container unsuitable for further use. Dispose of the container in accordance with provincial requirements. For information on the disposal of unused, unwanted product, and the cleanup of spills, contact the regional office of Conservation and Protection, Environment Canada.

TOXICOLOGY: Lindane is an organo-chlorine insecticide. Barbiturates may be given to control convulsions. Oxygen may be indicated. Avoid use of morphine and adrenaline.

NOTICE TO USER: This control product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products Act to use a control product under unsafe conditions.

LIMITATION OF WARRANTY: Seller's guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

FORESTRY PESTS: Logs (Balsam Fir, Pine, Spruce) - controls Sawyer Beetles, Ambrosia Beetle (Pinhole Borer), Pine Bark Beetles. Prepare 0.4% solution by using 1 litre in 25 litres of water. During May or June, spray cut logs to run-off. Ensure thorough coverage. Repeat if necessary. Do not put logs into water within 6 weeks after treatment.

CHRISTMAS TREES: (Austrian, Red and Scotch Pine) - controls cereal leaf beetle in overwintering stage. Quarantine treatment Ontario. All trees to be shipped from quarantined counties in Ontario to points outside restricted area in Canada and The United States must be treated with Lindane (or other product registered for this use) under the supervision of Plant Health Directorate, Agriculture Canada and must be provided with a quarantine movement certificate issued by Plant Health Directorate.

SPRAY: Mix 6 litres in 1,000 litres of water. Before cutting, spray trees to the point of run-off from two sides, at a pressure of about 2100 kPa. Thorough coverage is essential. A 1 1/2 to 2 metre tree requires about 2 litres for good coverage.

NURSERY STOCK, ORNAMENTAL (OUTDOORS) - Pine Bark Beetles - spray solution of 5 litres in 1,000 litres of Sanex Base Oil on soil and base of tree trunk.

INDOORS - controls ants, bees, booklice, carpet beetles, clothes moth, cockroaches, crickets, Drosophila, earwigs, fleas, gnats, hornets, house fly, larder beetle, meal moths, millipedes,

mosquitoes, silverfish, sowbugs, spiders, springtails, stable fly, wasps.

IN DWELLINGS - for residual, apply solution of 1 litre in 19 litres of water as a spot treatment to floors, walls and other infested areas. For clothes moths, treat surfaces of closets and storage chests. Do not use in cupboards where food is stored. Remove or cover all food before application. Do not use as a space spray. Remove fish and pets before spraying. After spraying, clean surfaces which may contact food.

IN FARM BUILDINGS (Including Barns, Creameries, Stables), FOOD PROCESSING PLANTS, WAREHOUSES - treat as above, in food processing plants, do not apply while plant is in operation. Any treated surfaces that may contact food should be washed and rinsed thoroughly. Do not use in milk processing rooms, milking parlours, or dairy barns. Do not contaminate food, feed utensils or areas within reach of animals. Do not use as a space spray.

For BEES, WASPS, HORNETS, indoors and outdoor surfaces of buildings, apply as a coarse wet 1 spray to nest, preferably at night while inactive.

BEEF CATTLE - prepare a solution of 3 to 6 litres in 1,000 litres of water. For hornfly, apply to back and shoulders at the rate of $1\ 1/4$ to $2\ 1/2$ litres per animal. For lice and ticks, apply as an overall spray using 4 to 9 litres per animal. For lice, repeat in 14 to 18 days.

GOAT (non-lactating), HORSE, SHEEP - Prepare a solution of 3-5 litres in 1,000 litres of water, apply as an overall spray to control fleas, lice, ticks using 2 to 5 litres per animal. For lice, repeat in 14 to 18 days.

HOGS - Prepare a solution of 3-5 litres in 1000 litres of water, apply 1 to 2 litres against fleas, lice, mange, mites, ticks.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.