

97.12.23

FURADAN (R) 5 GRANULAR

SYSTEMIC INSECTICIDE

Contains Carbofuran

COMMERCIAL

NET CONTENTS: _____KILOGRAMS

GUARANTEE: Carbofuran.....5%

REGISTRATION NO. 10666
PEST CONTROL PRODUCTS ACT

DANGER POISON

KEEP OUT OF REACH OF CHILDREN
(See Rear Panel for Additional Warning Statements)

READ THE LABEL BEFORE USING

FMC CORPORATION
1735 Market Street
Philadelphia, Pennsylvania
19103
USA

NOTICE TO USER: This control product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products Act to use a control product under unsafe conditions.

DIRECTIONS FOR USE

Read entire label and follow directions closely. Do not handle this product with bare hands.

| CROP | INSECT | DOSAGE - FURADAN 5 GRANULES | REMARKS |
|------|--------|--------------------------------|---------|
|------|--------|--------------------------------|---------|

| | | | |
|--------------------------------------|---------------------|--|--|
| <p>Canola (Rapeseed) and Mustard</p> | <p>Flea beetles</p> | <p>4.5 kg/ha to be increased to 5.5 kg in case heavy infestation is expected</p> | <p>For maximum flea beetle control apply a mixture of FURADAN 5 Granules and seed. Harrowing after seeding causes loss of efficacy. Mix 4.5 kg (in case of need 5.5 kg) of FURADAN 5 Granules with the seed to be sown per hectare. Example: If 5.5 kg of seed is sown per hectare, mix with 5.5 kg of FURADAN (total weight 11.00 kg). The FURADAN granules and seed should be thoroughly mixed using a customary mechanical method.</p> <p>BEFORE SEEDING, ADJUST THE SEEDING EQUIPMENT, SO THAT THE REQUIRED WEIGHT OF SEED AND FURADAN IS SOWN PER HECTARE. LOCATE AND BURY ANY SPILLS SUCH AS THOSE WHICH MAY OCCUR DURING CALIBRATION OR WHEN THE SEEDER IS LIFTED TO MAKE TURNS AT FIELD ENDS. ENSURE THAT THE SEED AND FURADAN GRANULES ARE COMPLETELY COVERED WITH SOIL IN ORDER TO MINIMIZE EXPOSURE TO BIRDS AND WILDLIFE.</p> <p>TO MINIMIZE GRANULE EXPOSURE TO BIRDS AND WILDLIFE, DO NOT BROADCAST SEED OR GRANULES ON THE SOIL SURFACE (WHETHER OR NOT FOLLOWED BY HARROWING) OR USE DOUBLE DISC PRESS DRILLS OR ANY DISCER TYPE SEEDERS.</p> <p>Check fields shortly after canola (rapeseed) emerges. Under certain conditions, a follow-up foliar treatment may be necessary. FURADAN 480 Flowable and GUTHION (R) are insecticides which may be used. If seed decay, seedling blight, and</p> |
|--------------------------------------|---------------------|--|--|

PRECAUTIONS:

DANGER: Poisonous by swallowing or inhalation. Do not breath dust. Do not get in eyes. Wear goggles when exposed to dust. In case breathing of dust during handling cannot be avoided, use a respirator of a type suitable for carbofuran protection such as American Optical Respirator 6500 (filter-cartridges R58) or Willson Agritox Respirator (Cartridge R21, filter R15). Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals. Do not contaminate feed or foodstuffs. Keep out of areas inhabited by fish, birds and wildlife, as this product is highly toxic to such animals.

To reduce potential wildlife hazard, any FURADAN 5 Granules left on the soil surface due to accidental spills or when the seeder is lifted to make turns should be covered with soil.

POISON

Call a doctor in case of accident

KEEP OUT OF REACH OF CHILDREN

FIRST AID

IF SWALLOWED, vomiting should be induced. Administer water freely and induce vomiting by giving one dose (15 mL) of syrup of ipecac. If vomiting does not occur within 10 to 20 minutes, administer second dose. If syrup of ipecac is not available, induce vomiting by sticking finger down throat. Repeat until vomit fluid is clear. The patient should be lying down with the head below the foot level and facing down or to one side. Professional medical assistance should be secured immediately. DO NOT INDUCE VOMITING TO AN UNCONSCIOUS PERSON OR TO PERSONS IN A CONVULSIVE STATE. IF INHALED, have victim lie down and remain quiet. IF ON SKIN, remove contaminated clothing and immediately wash skin with soap and water. IF IN EYES, flush eyes immediately with plenty of water for at least 15 minutes. Call a doctor immediately.

TOXICOLOGICAL INFORMATION: Symptoms of poisoning are headache, light-headedness, weakness, nausea, constriction of pupils, blurred vision, cramps, salivation, diarrhea, vomiting. Carbofuran is a reversible cholinesterase inhibitor. Give atropine 2 mg intramuscularly. If in eyes, instill one drop of homatropine. Do not use oximes such as 2-PAM..

DISPOSAL:

20 KG SIZE: Completely empty the contents into the pesticide hopper. Make the empty container unsuitable for further use. Do not reuse container. Dispose of the container in accordance with provincial requirements but DO NOT BURN.

500 KG SIZE: After use, return the container to the point of purchase or designated locations. This container must only be refilled with FURADAN 5 GRANULES. Do not reuse container for any other purpose. Prior to refilling inspect thoroughly for damage such as crack, punctures, abrasions, and damaged or worn out threads on closure devices. Check for leaks after refilling and before transportation. Do not refill or transport damaged or leaking containers. If the container is not being refilled, return to point of purchase.

For information on the disposal of unused, unwanted product and the cleanup of spills, contact the Provincial Regulatory Agency or the Manufacturer.

NOTICE TO BUYER

Seller's guarantee shall be limited to the terms set out on the label and, subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))