

96.06.07

CLEAN CROP(R)

STAMPEDE*
360

HERBICIDE

COMMERCIAL

WARNING

POISON CORROSIVE

READ THE LABEL BEFORE USING

GUARANTEE:

Propanil..... 360 g/L

REGISTRATION NO. 17383
PEST CONTROL PRODUCTS ACT

PRECAUTIONS

Causes severe skin and eye irritation. Avoid contact with skin and eyes. Wear protective goggles when handling this concentrate. Do not breathe spray mist. Avoid swallowing. Do not store near feed or food products. Keep out of reach of children. This product is toxic to fish. Keep out of streams, ponds and irrigation waters. Do not contaminate water by cleaning of equipment or disposal of waste or rinsate. Avoid spray drift off area of application.

FIRST AID

In case of contact with skin, wash thoroughly with soap and water. In case of contact with eyes, flush with plenty of water for at least 15 minutes and get medical attention or contact a poison control centre. If taken internally, dilute by giving 2 glasses of water to drink and call a physician. Never give anything by mouth to an unconscious person.

UNITED AGRI PRODUCTS
R.R. #2
Dorchester, Ontario
N0L 1G5

Reg. T.M. of Rohm & Haas Company

NET: 11.36L

DIRECTIONS FOR USE

STAMPEDE 360 is a selective postemergence herbicide for the control of green and yellow foxtail (wild millet, pigeongrass) and certain broadleaf weeds in spring wheat (including durum), barley, oats, canary grass (canary seed), and flax. In common with most herbicides, STAMPEDE 360 is more effective on activity growing weeds. For best results tank-mix STAMPEDE 360 with MCPA ester (or with 2,4-D Amine or 2,4-D Ester or GLEAN 75 DF on wheat and durum) and apply when weeds are small and growing conditions are favourable for vigorous growth:

WEEDS CONTROLLED

A. STAMPEDE 360 tank-mix with MCPA Ester (on wheat, barley, oats, canary grass (canary seed), and flax).

Green foxtail (wild millet, pigeongrass, <i>Setaria viridis</i>)	
Yellow foxtail (wild millet, pigeongrass, <i>setaria lutescens</i>)	
Wild buckwheat	Tartary buckwheat
Smartweed	Lady's-thumb
Redroot pigweed	Lamb's-quarters
Stinkweed*	Shepherd's-purse*
Wild mustard	Flixweed*
Volunteer rapeseed	Kochia*
Bluebur*	*in seedling or rosette stage

B. STAMPEDE 360 tank-mix with 2,4-D Amine or 2,4-D Ester (on wheat only including durum):

Weeds listed in A plus:

Russian thistle	Sweet clover
Cocklebur	Burdock (seedling)
Narrow-leaved hawk's beard (fall rosette and spring seedling)	Goat's beard (seedling)
Plantain	Prickly lettuce
Russian pigweed	Annual sunflower
	Wild radish

Note: The 400 g of 2,4-D per hectare rate in the tank-mix may not provide adequate control of the above weed species. Refer to the 2,4-D label.

C. STAMPEDE 360 tank-mix with 15 g/ha (low rate) of GLEAN 75 DF (on wheat, durum and barley varieties Argyle, Bedford and Klages)

Green Foxtail
(wildmillet, pigeongrass)
Yellow foxtail
(wildmillet, pigeongrass)
Cow cockle
Flixweed
Green smartweed
Hemp-Nettle
Lady's-thumb
Lamb's-quarters

Redroot pigweed
Shepherd's-purse
Stinkweed
Tartary buckwheat
Volunteer rapeseed
Wild buckwheat
Wild mustard

D. STAMPEDE 360 tank-mix with 30 g/ha (high rate) of GLEAN 75 DF
(on wheat, durum and barley varieties Argyle, Bedford, Klages)

Weeds listed in C plus:

Chickweed
Kochia
Cleavers
Russian thistle
Canada thistle*

*For seasonal control only. Must be sprayed in the flower bud stage.

APPLICATION WEED STAGE

Apply STAMPEDE 360 tank-mix with MCPA or 2,4-D Amine, 2,4-D Ester or GLEAN 75 DF when the majority of green foxtail plants are in the 3 leaf stage and less than 2.5 centimetres in height. For best yield response, it is important to control the flush of green foxtail that emerges with the crop. The effectiveness of the tank-mix in controlling green foxtail declines rapidly as the 5th leaf begins to emerge. Under prolonged dry conditions apply STAMPEDE 360 tank mixes when the green foxtail is in the 2-3 leaf stage. Susceptible broadleaf weeds in the 1-4 leaf stage are also controlled with a tank-mix. The treatment will not control green and yellow foxtail and broadleaf weeds that emerge through the soil after application.

CROP STAGE AND RATE OF APPLICATION:

- A) STAMPEDE 360 tank-mix with MCPA Ester . This tank-mix can be applied on wheat, durum, barley or oats at the 2-5 leaf stage and to flax from 5 cm to 12.5 cm in height. Apply 2.75 litres of STAMPEDE 360 per hectare plus 275 grams of MCPA Ester acid equivalent per hectare (equivalent to 0.55 litres of a formulation containing 500 grams of MCPA acid equivalent per litre).
- B) STAMPEDE 360 tank-mix with 2,4-D Amine or 2,4-D Ester. This tank-mix can be applied to only spring wheat and durum when

the crop is in the 3-5 leaf stage. Apply 2.75 litres of STAMPEDE 360 per hectare plus 600 grams of 2,4-D Amine acid equivalent per hectare (equivalent to 1.2 litres of a formulation containing 500 grams 2,4-D Amine acid equivalent per litre). Apply 2.75 litres of STAMPEDE 360 per hectare plus 400 to 600 grams of 2,4-D Ester acid equivalent per hectare on spring wheat only (equivalent 0.66 to 1.0 litres of formulation containing 600 grams 2,4-D Ester acid equivalent per litres). Apply a maximum of 400 grams of 2,4-D Ester acid equivalent per hectare on durum wheat only.

- C) STAMPEDE 360 and GLEAN 75 DF tank mix. This tank-mix can be applied on wheat including durum and barley when the crop is in the 2-5 leaf stage. Apply 2.75 litres of STAMPEDE 360 per hectare plus 15 g/ha of GLEAN 75 DF for control of the weeds listed above in part C and 30 g/ha of GLEAN 75 DF for control of weeds listed above in part D.

MIXING

STAMPEDE 360 is an emulsifiable concentrate and mixes readily with water. It is recommended that the water in the spray tank be 10 degrees C or warmer when the STAMPEDE 360 is added. When mixing, add the MCPA 2,4-D or GLEAN 75 DF first to a half-filled spray tank; then STAMPEDE 360. Maintain agitation during filling. Do not let mixture stand without agitation.

EQUIPMENT

Apply STAMPEDE 360 with a properly calibrated low pressure ground sprayer employing 50 mesh screens (both nozzles and filter screens). Use 100 litres of water per ha at 275 kPa. Use flat fan nozzles (such as Tee Jet 6502, or 8002, LF-2-65' or LF-2-80', etc.). Drain and flush spray tank and lines thoroughly at the end of each day's spraying and before switching to or from another agricultural chemical.

Floater equipped with float-jet nozzles are satisfactory for STAMPEDE 360 application. A minimum of 150 litres of water per hectare is recommended to overcome the coarser water droplets produced. The nozzles should be spaced such that complete overlap of spray patterns occur. Applications by aircraft or spray coupe are not recommended.

CROP TOLERANCE

STAMPEDE 360 may cause temporary crop yellowing or tip browning to wheat, barley, oats and flax 2-5 days after application. The effects are temporary and usually disappear 10-14 days after application. New leaves will have the normal green colour. Application of STAMPEDE 360 under adverse growing conditions may result in greater crop injury and slower recovery. STAMPEDE 360 should not be applied on crops under growth stress caused by drought, flooding, excess soil salts, hail damage, lack of

fertilizer, disease (foliar or root) or extreme temperatures. (See WEATHER for further information on crop tolerance).

WEATHER

Rain: Light rain one hour after application does not reduce yield control. A heavy rain of 25 mm or more, within 4 hours of application may reduce control.

Soil Moisture: Weed control with STAMPEDE 360 is strongly influenced by soil moisture. Weed control and crop tolerance is best when adequate moisture is available for active weed and crop growth. The activity of STAMPEDE 360 is reduced substantially during periods of prolonged dry weather. Under prolonged dry conditions STAMPEDE 360 should be applied earlier, when the green foxtail is in the 2-3 leaf stage. STAMPEDE 360 is not recommended for use when soil moisture levels are deeper than 5 cm.

Humidity and Temperature: STAMPEDE 360 is most effective when relative humidity is high and daily temperatures exceed 21 degrees C. If weather is hot, dry and relative humidity low, restrict spraying to early morning or evening. Do not spray STAMPEDE 360 when daily maximum temperature will remain below 10 degrees C or when they are expected to exceed 30 degrees C.

Frost: Avoid applying STAMPEDE 360 if frost is expected within 24 hours and do not apply STAMPEDE 360 to a crop damaged by frost until fully recovered.

Wind: Avoid applying STAMPEDE 360 to crops damaged as the result of high wind until the crop is fully recovered.

COMPATIBILITY WITH OTHER CHEMICALS

Always follow the label recommendations on agricultural chemicals mixed with STAMPEDE 360.

FOLIAR APPLIED CHEMICALS

Herbicides

STAMPEDE 360 can be tank mixed MCPA Ester, 2,4-D Ester or Amine and GLEAN 75 DF herbicides. If another herbicide is required for example, a post-emergence wild oat herbicide, a 3 day interval should be allowed between the application of STAMPEDE 360 and the other herbicide.

Insecticides

Severe injury or crop plants may result from a tank-mix or separate applications of STAMPEDE 360 herbicide and certain insecticides in the same crop year, i.e. SEVIN (carbaryl), Parathion methyl or GUTHION.

If a foliar insecticide is necessary, DECIS (deltamethrin) may be

applied at anytime before or after an application of STAMPEDE 360 or as a tank mix with STAMPEDE 360. Consult the DECIS label for application rate and other instructions.

FURADAN (carbofuran) may be applied on wheat 5 days or longer after an application of STAMPEDE 360 or 10 days after an application of STAMPEDE 360 on barley. CYGON (dimethoate) (LAGON) or MALATHION may be applied 14 days after an application of STAMPEDE 360. No other insecticides are registered for foliar use in the same year as STAMPEDE 360 is applied.

Fertilizers and Spray Adjuvants

Do not tank mix STAMPEDE 360 with foliar applied fertilizers or spray adjuvants.

SOIL APPLIED CHEMICALS

HERBICIDES

STAMPEDE 360 can be applied to crops grown in soil treated in the fall or spring with AVADEX BW (triallate) preplant incorporated for wild oat control. DO NOT APPLY STAMPEDE 360 TO CROP GROWN IN FIELDS IN WHICH ATRAZINE HAS BEEN APPLIED IN EITHER OF THE PREVIOUS TWO YEARS.

INSECTICIDES

Do not spray crops with STAMPEDE 360 if the field was treated with soil applied systemic organophosphorous insecticides (DISYSTON, THIMET, COUNTER, etc.) in the same or previous crop year.

SEED TREATMENT

Crops grown from seed protected with dual-purpose (insecticide/fungicide) seed dressings may be treated with STAMPEDE 360.

CAUTION: Do not graze the treated crops or cut for hay; sufficient data are not available to support such use.

PRECAUTIONS

Causes severe skin and eye irritation. Avoid contact with skin and eyes. Wear protective goggles when handling this concentrate. Do not breathe spray mist. Avoid swallowing. Do not store near feed or food products. Keep out of reach of children. This product is toxic to fish. Keep out of streams, ponds and irrigation waters. Do not contaminate water by cleaning and equipment or disposal of waste or rinsate. Avoid spray drift off area of application.

Thoroughly clean spray equipment before using for other purposes. Following cleaning procedure outlined for 2,4-D and MCPA Ester removal in provincial Weed Control Manuals.

FIRST AID

In case of contact with skin, wash thoroughly with soap and water. In case of contact with eyes, flush with plenty of water for at least 15 minutes and get medical attention or contact a poison control centre. If taken internally, dilute by giving 2 glasses of water to drink and call a physician or contact a poison control centre. Never give anything by mouth to an unconscious person.

TOXICOLOGY

Treat as above for skin and eye contact. If taken internally, emesis is recommended because of the toxicity of this product. However, because of the presence of petroleum hydrocarbons, careful gastric lavage may be indicated. Morphine or morphine derivatives and barbiturates should be avoided. Studies in laboratory animals indicate that excessive exposure may result in central nervous system and neuromuscular depression.

DISPOSAL

1. Rinse the emptied container thoroughly and add the rinsings to the spray mixture in the tank.
2. Follow provincial instructions for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.
4. Dispose of the container in accordance with provincial requirements.
5. For information of the disposal of unused, unwanted product and the cleaning up of spills, contact the provincial regulatory agency or the manufacturer.

NOTICE TO USER

This control product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products Act to use a control product under unsafe conditions.

NOTICE TO BUYER

Seller's guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this

service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))