

85.12.31

KEMSAN ABATE* 2G
GRANULAR INSECTICIDE
KEMSAN ABATE* 2G
INSECTICIDE GRANULAIRE

RESTRICTED/A USAGE RESTREINT

NET CONTENTS/CONTENU NET: 20 kg

CAUTION/ATTENTION: POISON

GUARANTEE/GARANTIE: TEMEPHOS...2%

REG.NO./NO.D'ENR. 13695 P.C.P. ACT/L.P.A.

READ THE LABEL BEFORE USING
LIRE L'ETIQUETTE AVANT L'EMPLOI

KemSan INC.
P.O. BOX 727
OAKVILLE, ONTARIO
L6J 5C1
TELEPHONE (416) 845-2271

FORMULATING A BETTER FUTURE FOR YOU

KemSan INC.
C.P. 4835
SAINT-LAURENT, QUEBEC
H4L 4Z5
TELEPHONE (514) 332-5744

VOUS OFFRE DES FORMULES DE SUCCES

NOTICE TO USER: This control product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products Act to use a control product under unsafe conditions.

NATURE OF RESTRICTION: This product is to be used only in the manner authorized; consult local authorities about use permits which may be required.

RESTRICTED USES USE LIMITATIONS: Remove all livestock during application.

TO CONTROL MOSQUITO LARVAE: Use ABATE 2-g Insecticide as directed below.

AREAS OF TREATMENT	RATE KG PER HECTARE
Standing water in shallow ponds, woodland pools, catch basins, etc. (relatively clean open water)...	2.75 to 5.50
Tidal water, marshes, swamps (water high in organic matter content)...	11
Highly polluted water.....	27.50

The higher rates should be used in areas known to have organophosphate resistant mosquitoes.

Apply uniformly by ground or aid equipment.

Repeat application as necessary.

AERIAL APPLICATION. Apply ABATE 2-G granular by means of fixed-wing aircraft or helicopter at the rate of 2.75 to 5.50 kg/ha to standing water, shallow ponds and woodlot ponds.

In marshlands, swamps and waters of high organic matter, apply ABATE 2-G granular at the rate of 11 kg/ha.

Apply product uniformly to those areas that require treatment only.

MOSQUITO LARVAE CONTROL IN FIRE AND FARM PONDS ONLY: For use in water wholly confined within the property of the user and where there is no outflow beyond the property limits.

Apply in the spring when larvae are just noticed in the water. Retreatment may be necessary in 6 to 8 weeks after initial treatment.

Apply ABATE 2-G granular at the rate of 2.75 to 5.50 kg/ha.

In waters high in organic matter content, apply at the rate of 11 kg/ha.

PRECAUTIONS: HARMFUL IF SWALLOWED, or absorbed through the skin. Do not get on skin, in eyes or on clothing. Wash thoroughly after handling. Avoid breathing dust. Do not contaminate food or feed products. Causes eye and skin irritation. Keep out of reach of children. Do not expose product to excessive heat or

direct sunlight. Do not apply to crops used for food, forage or pasture.

FIRST AID: In case of contact, flush skin or eyes with plenty of water. Call a physician immediately in all cases of suspected poisoning.

TOXICOLOGICAL INFORMATION: This product may cause cholinesterase inhibition. Atropine is antidotal. Pralidoxime chloride(2-PAM; PROTOPAM chloride) may be effective as an adjunct to atropine. Use according to label directions.

THIS PRODUCT IS TOXIC TO BIRDS. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not reuse empty container. Destroy it by burying with wastes or burning. Stay away from smoke and fumes.

NOTICE TO BUYER: Seller's guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

* Registered Trademark CYANAMID CANADA INC.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))