

**SNARE RODENTICIDE
BAIT PELLET
KILLS RATS AND MICE
Kills Warfarin Resistant Norway Rats**

Norway rat and house mice may consume a lethal dose in one feeding with first dead rodents appearing four or five days after treatment begins.

REG. NO. 23167 P.C.P. ACT
DOMESTIC

**WARNING
POISON**

GUARANTEE: Bromadiolone 0.005%
Warning, contains the allergen wheat.
Contains malathion at 0.01%, as a preservative.

READ THE LABEL BEFORE USING/KEEP OUT OF REACH OF CHILDREN

NET CONTENTS: 35 x 50 g (1.8 oz.)

Distributed by: CITADEL ANIMAL HEALTH, Cambridge, ON, N3C 2W4

USE LIMITATIONS: Bait MUST either be placed in tamper-resistant bait stations or in locations not accessible to children, pets or livestock. DO NOT place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food.

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN, PETS AND LIVESTOCK. May be harmful or fatal if swallowed or absorbed through the skin. Chemical-resistant gloves must be worn when handling product and when disposing of dead rodents, unconsumed bait and empty containers. Avoid contact with eyes, skin or clothing. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Wash skin thoroughly with soap and water after handling. Wash contaminated clothing, separately from other laundry, with soap and hot water before reuse. KEEP AWAY FROM FEED AND FOODSTUFFS. Do not expose treated bait in areas accessible to children, pets, domestic animals and wildlife, except when enclosed within a tamper-resistant bait box. If swallowed, this material may reduce clotting ability of the blood and cause bleeding. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food. Keep out of lakes, streams or ponds.

FIRST AID: If in eyes, flush with water for 15 minutes. If on skin, wash with soap and water. If swallowed, give patient one or two glasses of water and induce vomiting by touching back of throat with a finger or a blunt object; repeat until vomit fluid is clear. Do not give anything by mouth to an unconscious person. For all cases of human ingestion, immediately notify a physician or poison control centre. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention. If pet or livestock poisoning is suspected, immediately contact a veterinarian.

TOXICOLOGICAL INFORMATION: Vitamin K₁, in the form of intramuscular or subcutaneous injections, or by oral ingestion, is the suggested remedial treatment for anticoagulant poisoning. The severity of the case measured by establishing prolonged prothrombin times (P.T.) will determine appropriate therapy. Monitoring P.T. will indicate necessity of repeated treatments.

APPLICATION DIRECTIONS:

RATS: Place 100 to 500 grams of bait at intervals of 4.5 to 9 metres per placement. Maintain an uninterrupted supply of fresh bait for 10 days or until signs of rat activity cease.

HOUSE MICE: Place 7 to 14 grams of bait at intervals of 2.5 to 3.5 metres per placement. Larger placements (up to 100 grams) may be needed at points of high mouse activity. Maintain an uninterrupted supply of fresh bait for 15 days or until signs of mouse activity cease.

RATS/MICE: Replace contaminated or spoiled bait immediately. Collect and dispose of unconsumed bait and dead exposed animals according to local or provincial requirements. To prevent reinfestation, eliminate food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish bait as needed.

DIRECTIONS FOR USE: Use bait to control the Norway rat (*Rattus norvegicus*), Roof rat (*Rattus rattus*) and House mouse (*Mus musculus*) indoors and against outside walls of buildings.

SELECTION OF TREATMENT AREAS: Determine areas where rats and/or mice will most likely find and consume the bait. Generally, these areas are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been observed. Remove as much food as possible.

STORAGE: Store in a cool, dry place away from other chemicals and food or feed. Store product not in use, in original container, in a secure location inaccessible to children and non-target animals.

DISPOSAL: Do not reuse empty container. Dispose of empty container by placing in household garbage.

RODENT DISPOSAL: Dispose of dead rodents in garbage or by burying.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.