

26-MAR-2007

25-FEB-2008 Notification - Change in Registrant Name

(THROW BAG)

**WILSON
BROMONE[®]
RODENTICIDE MEAL BAIT**

DOMESTIC

REG. NO. 16700 P.C.P. ACT

GUARANTEE: BROMADIOLONE 0.005%

Contains malathion at 0.0096% and calcium propionate at 0.3% as preservatives

WARNING



POISON

READ THE LABEL AND OUTER CARTON BEFORE USING
KEEP OUT OF REACH OF CHILDREN

50 g

Notification Change

~~Nu-Gro PR Inc.~~
10 Craig St.
Brantford, ON
Canada N3R 7J1
519-757-0077

Agrium Advanced Technologies RP Inc.

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN, PETS AND LIVESTOCK.
Before using, see outer carton for full PRECAUTIONS regarding placement and storage.

FIRST AID: **If in eyes,** flush with water for 15 minutes. **If on skin,** wash with soap and water. **If swallowed,** give patient one or two glasses of water and INDUCE VOMITING by touching back of the throat with a finger or blunt object; repeat until vomit fluid is clear. Never give anything by mouth to an unconscious person. For all cases of human ingestion, immediately notify a physician or poison control centre. If pet or livestock poisoning is suspected, immediately contact a veterinarian. Take container, label or product name and P.C.P. Reg. No. with you when seeking medical attention.

DISPOSAL: Wrap container and discard in household garbage. Do not reuse empty container.
RODENT DISPOSAL: Dispose of dead rodents in garbage or by burying.

(CARTON)

26-MAR-2007

**WILSON
BROMONE®
RODENTICIDE MEAL BAIT**

One Feeding Kills Rats And Mice

DOMESTIC

REG. NO. 16700 P.C.P. ACT

GUARANTEE: BROMADIOLONE 0.005%
Contains malathion at 0.0096% and calcium propionate at 0.3% as preservatives

WARNING



POISON

READ THE LABEL BEFORE USING
KEEP OUT OF REACH OF CHILDREN

1 kg (20 x 50 g bait bags)

Notification Change

~~Nu-Gro PR Inc.~~
10 Craig St.
Brantford, ON
Canada N3R 7J1
519-757-007

Agrium Advanced Technologies RP Inc.

USE LIMITATIONS

Bait **MUST** either be placed in tamper-resistant bait stations or in locations not accessible to children, pets or livestock. **DO NOT** place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food.

DIRECTIONS FOR USE (indoors and against the outside walls of buildings): For the control of rats and mice.

Rats: Place 120 g of bait at 3 to 4 metre intervals. Maintain an uninterrupted supply of fresh bait for at least 10 days or until signs of rat activity cease.

Mice: Place at least 10-15 g of bait at intervals of 2.5 to 3.5 metres per placement. Larger placements (30-60 g) may be needed at points of high mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 days or until signs of mouse activity cease.

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN, PETS AND LIVESTOCK.

May be harmful or fatal if swallowed or absorbed through the skin. Do not open pre-measured place packs. Rubber gloves must be worn when disposing of dead rodents, unconsumed bait and empty containers. Avoid contact with eyes, skin or clothing. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Wash skin thoroughly with soap and water after handling. Wash contaminated clothing, separately from other laundry, with soap and hot water before reuse. **KEEP AWAY FROM FEED AND FOODSTUFFS.** This product is toxic to fish and wildlife. Keep out of lakes, streams or ponds.

FIRST AID: If in eyes, flush with water for 15 minutes. **If on skin,** wash with soap and water. **If swallowed,** give patient one or two glasses of water and **INDUCE VOMITING** by touching back of the throat with a finger or blunt object; repeat until vomit fluid is clear. Do not give anything by mouth to an unconscious person. For all cases of human ingestion, immediately notify a physician or Poison Control Centre. If pet or livestock poisoning is suspected, immediately contact a veterinarian. Take container, label or product name and P.C.P. Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION: This product contains an anti-coagulant chemical which, if taken accidentally by humans or pets, may reduce the clotting ability of the blood and haemorrhage may result. Vitamin K₁ in the form of intramuscular or subcutaneous injections, or by oral ingestion are suggested remedial treatments for anticoagulant poisoning. The severity of the case, measured by establishing prolonged prothrombin times (P.T.), will determine appropriate therapy. Monitoring P.T. will indicate the necessity of repeated treatments.

STORAGE:

Store in a cool, dry place away from other chemicals and food or feed. Store product not in use, in original container, in a secure location inaccessible to children and non-target animals.

DISPOSAL: Wrap container and discard in household garbage. Do not reuse empty container.
RODENT DISPOSAL: Dispose of dead rodents in garbage or by burying.

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.