

89.06.09

IPCO (R)

COBUTOX 400

EMULSIFIABLE CONCENTRATE HERBICIDE

10 LITRES

AGRICULTURAL

FOR WEED CONTROL IN SEEDLING ALFALFA,
BIRDSFOOT TREFOIL AND PASTURES

READ THE LABEL AND ENCLOSED PAMPHLET BEFORE USING

GUARANTEE: 2,4-DB 400 g/L (Present as Ester)

REGISTRATION NO: 20315
PEST CONTROL PRODUCTS ACT

WARNING POISON

Interprovincial Cooperative Limited,
P.O. Box 1050, Saskatoon, Saskatchewan S7K 3M9
Calgary, Regina, Winnipeg, Mississauga, Montreal, Moncton

(R) T.M. Reg'd.

Dear Customer:

This product has a new label format:

1. It is easier to read. (Larger type, better layout and graphics)
2. It contains safety information which, if followed, will reduce your exposure and protect the environment.
3. The information you need is easier to find.
4. Responsible care is the responsibility of all of us, manufacturer, distributor, dealer and user. Help us to establish more public trust in the use of pesticides by handling all products as safely as possible.

Steps to Safe Use:

1. Wear rubber gloves and coveralls before handling any crop

- protection product.
2. Fill all spray tanks away from wells, sloughs, creeks or any other water way.
 3. Use a nurse tank or long hose to fill your spray tank.
 4. Wash off any spills immediately. Wash daily all clothing worn while filling and spraying. Use a separate wash for contaminated clothing.
 5. Do not spray when wind exceeds 15 km/h.
 6. Store all chemicals in a safe place under lock and key.
 7. Always store chemicals in the original container.

LABEL INDEX Cobutox 400	
Seedling Alfalfa and Birdsfoot Trefoil (direct or underseeded)	GO TO SECTION 1
Seedling Clover (direct or underseeded)	GO TO SECTION 2
Cereals (Wheat, Oats, Barley)	GO TO SECTION 3
Field Corn	GO TO SECTION 4
Pastures Containing Forage Legumes	GO TO SECTION 5
Weeds Controlled and Rate	GO TO SECTION 6
Tank Mix with MCPA (To improve wild mustard control in seedling alfalfa)	GO TO SECTION 7
Filling Instructions	GO TO SECTION 8
Precautions	GO TO SECTION 9
First Aid	GO TO SECTION 10
Rinsing and Disposal	GO TO SECTION 11

SECTION 1	
Crop:	SEEDLING ALFALFA AND BIRDSFOOT TREFOIL (direct or underseeded) NOTE: Do Not Apply if Crop is to be used for Seed Production Later in the Season.

Treatment Stage:	Treat from 1st to 4th trifoliolate leaf stage.
GO TO SECTION 6	
SECTION 2	
Crop:	SEEDLING CLOVERS (direct or underseeded) NOTE: Do Not Spray Sweet Clover. Red and Alsike Clovers may be damaged.
Treatment Stage:	Treat as soon as possible after the first trifoliolate leaf appears.
GO TO SECTION 6	
SECTION 3	
Crop:	CEREALS (WHEAT, OATS, BARLEY)
Treatment Stage:	Apply after the crop has 5 fully expanded leaves but before the early flag leaf stage. Oats will be damaged if treated before 5 leaf stage.
GO TO SECTION 6	
SECTION 4	
Crop:	FIELD CORN
Treatment Stage:	Apply after crop is at least 40 cm high but before tasselling. Use drop nozzles to avoid spraying corn leaves.
GO TO SECTION 6	
SECTION 5	
Crop:	PASTURES CONTAINING FORAGE LEGUMES

Treatment Stage:	Treat after cutting or grazing when growth is not above 7.5 cm in height. Damage may occur if crop is sprayed beyond this stage.
GO TO SECTION 6	

SECTION 6		
Annual Weeds Controlled:	Rate:	One Jug Treats
Wormseed Mustard, Wild Mustard, Ball Mustard, Stinkweed, Ragweed, Redroot Pigweed, Shepherd's Purse, Lamb's-quarters.	2.75 L/ha	3.6 ha (9 acres)
Oakleaved Goose Foot, Wild Buckwheat, Wild Radish (NOTE: Best control occurs when weeds sprayed in the 1-2 leaf stage.)	3.5 L/ha	3.85 ha (7 acres)
Lady's-thumb, Green Smartweed. Seedlings stunted only, but competitive ability is reduced.	4.25 L/ha	2.35 ha (5.8 acres)

SECTION 7	
Tank Mix with MCPA:	To improve wild mustard and narrow leafed hawksbeard control in seeding Alfalfa.
Treatment Stage;	Treat mustard beyond four leaf stage. Apply Cobutox alone on mustard at 4 leaf stage or less.
Rate:	Cobutox 2 L/ha plus 5 ha (12.5 acres) +MCPA Amine 500 70 mL/ha 140 ha (350 acres)
GO TO SECTION 8	

SECTION 8

FILLING SPRAY TANK

1. Fill tank 1/2 full with water.
2. Add Cobutox
3. When MCPA Amine 500 is used, add and mix before adding Cobutox. 4. Complete filling of tank with water.
5. Use 150-200 lites of water per hectare.
6. Do not exceed 275 kPa pressure.

GO TO SECTION 9

SECTION 9

PRECAUTIONS:

1. Keep out of reach of children.
2. Avoid contact with eyes, skin and clothing. May cause skin irritation.
3. Do not breathe spray mist or vapors. Avoid spray or vapor drifts onto susceptible plants such as grapes, tomatoes, tobacco, sugar beets, sunflowers, beans, turnips, cauliflower, cabbage, alfalfa, clover, peas, potatoes, ornamental trees, shrubs or flowers, fruit trees, lawns, most vegetable crops or other desirable vegetation.
4. Do not spray when there is any danger of wind or on exceptionally hot days over 27 deg. C.
5. Drift from spray may be reduced by using high volume sprays under low pressure, coarse sprays, and drop nozzles. Use special care in airplane application where danger from drift is greater.
6. For aerial application exercise special care:
 - 1) Do not spray in winds exceeding 8 km per hour.
 - 2) Do not spray in dead calm near sensitive plants. A cloud of suspended droplets may drift onto sensitive plants when the wind comes up.
 - 3) Spray only when the wind is blowing away from a sensitive crop, shelter belt or garden.
7. Do not contaminate any body of water. Avoid contamination of foods.
8. Do not use in a greenhouse.

9. Keep in original container during storage.
10. Do not store near or in same room as seeds, feeds, fertilizers or pesticides.
11. Keep away from fire or open flame and other sources of heat.
12. Clean spray equipment thoroughly after use. Do not use spray equipment for other purposes. DO not re-use empty containers.

GO TO SECTION 10

SECTION 10

FIRST AID:

1. In case of poisoning, call a physician immediately.
2. If on skin, remove contaminated clothing and wash skin thoroughly with soap and water.
3. If in eyes, flush with water immediately and continue for 5-10 minutes. Obtain medical attention.
4. If swallowed, do not induce vomiting, but rush patient to nearest hospital or doctor's office. Take the pesticide label with you.

GO TO SECTION 11

SECTION 11

Disposal Information:

Dear Customer: It is your responsibility to:

1. Triple rinse or jetrinse each container and add each rinse to spray solution. A rinsed container reduces the potential for environmental contamination to almost ZERO.
2. Dispose of the properly rinsed container in a manner approved in your province. By properly rinsing empty containers, you will help to reduce the opposition to the use of crop protection products.

NOTICE TO USER:

This control product is to be used only in accordance with the directions on this label. It is an offense under the Pest Control Products Act to use a control product under unsafe conditions.

NOTICE TO BUYER:

Seller's guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use of handling of this product and accepts the product on that condition.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))