

**05-JUL-2004 Notification received of a label change. See below for this change.**

17-MAR-1989

CLEAN CROP®

ZINEB 80W

WETTABLE POWDER

FUNGICIDE

COMMERCIAL

CAUTION....POISON

READ THE LABEL BEFORE USING

GUARANTEE: Zineb.....80%

REGISTRATION. No. 14562

PEST CONTROL PRODUCTS ACT

NET 10 kg

Notification Change

**UNITED AGRI PRODUCTS CANADA INC.□**

**789 DONNYBROOK DRIVE□**

**DORCHESTER ON N0L 1G5**

~~UNITED AGRI PRODUCTS~~

~~P.O. BOX 22116~~

~~LONDON, ONTARIO~~

~~N6C 4N0~~

DIRECTIONS

Fill spray tank 1/4 to 1/2 full with water, start agitators and slowly add the Zineb 80W. When mixed well, fill tank with water. Spray to provide thorough coverage of all plant surfaces.

NOTE: Numbers following crops listed below indicate the minimum number of days applications can be made up to harvest.

Crop Pest	Method of Application	Rate per 1000 Litres
POTATO (1) Early Blight, Late Blight	Commence application when plants are 10-15 cm high and repeat at 7 to 10 days intervals throughout the growing season. Use the lower rates early in the season, and increase rates as vines increase in size.	2-3 Kg
BROCCOLI, BRUSSEL SPROUTS, CABBAGE, CAULIFLOWER (7) Black Leaf Spot, Downy Mildew, Gray Leaf Spot	Apply when the disease first appears and repeat at 5 to 10 day intervals as necessary.	2 Kg
CUCUMBERS, MELON, PUMPKIN, SQUASH (5) Alternaria Leaf Spot, Anthracnose, Black Rot, Downy Mildew, Septoria Leaf Spot	Apply when plants begin to vine or when disease threatens and repeat at 5 to 10 day intervals as required.	2-2.5 Kg
TOMATOES (5) Anthracnose, Early Blight, Late Blight, Gray and Septoria Leaf Spot	Commence application after first fruit is well formed and repeat applications at 7 to 10 day intervals.	2-3.0 Kg
CARROT (7) Alternaria Blight, Carcospora Blight	Apply when disease first appears and repeat every 7 to 10 days as required.	2.75 Kg
CELERY (14) Early Blight, Late Blight	Apply to the plants in the seedbed and repeat applications in the field at weekly intervals.	2-3 Kg
LETTUCE (10) Downy Mildew	Apply when disease first appears and repeat every 7 to 10 days as required.	2-2.5 Kg

ONION (7) Botrytis Leaf Blight, Downy Mildew, Purple Blotch	Apply when disease first appears and repeat at 7 to 10 day intervals.	2-2.75 Kg
APPLE (14) Cedar Apple Rust, Fly Speck, Scab, Sooty Blotch	Apply in the 5th and 6th cover sprays. Make 1 more application 2 weeks after 6th cover spray.	1-1.25 Kg
PEAR (7) Sooty Blotch	Apply in the 2nd cover spray and repeat 2 weeks later.	2 Kg
TOBACCO Blue Mold	Apply 24-62 litres of spray per 100 m <sup>2</sup> . Apply when seedlings are small and before disease appears. Repeat applications every 3 to 5 days until transplanting time. Use lower rates on very small plants.	95-250 g
HOPS (24) Downy Mildew	Apply every 7 to 10 days as required.	1.5 to 2 Kg
ORNAMENTALS: Septoria Leaf Spot of Chrysanthemums, Rust of Snapdragons. Apply spray every 3-5 days as needed using 1.50 to 2.0 Kg/1000 L.		

#### CAUTION

Keep away from food and feed. Wash thoroughly after handling and before eating, drinking or smoking. Wash contaminated clothing with soap and hot water before reuse. Do not contaminate any body of water. May be harmful if inhaled or swallowed. May cause irritation of eyes, nose, throat or skin. Avoid breathing dust or spray mist. Avoid contact with skin, eyes and clothing. Keep away from fire or open flame. Never allow Zineb to become wet in storage as the fungicide effectiveness may be impaired and flammable vapours may form. Keep container closed when not in use. KEEP OUT OF REACH OF CHILDREN.

#### FIRST AID

If on skin, remove contaminated clothing and wash skin with soap and water. If in eyes, flush with clear water for 5 to 10 minutes and

obtain medical attention. If swallowed, induce vomiting by administering an ametic such as salty or soapy water. Repeat until vomit fluid is clear. Obtain prompt medical attention.

#### DISPOSAL

1. Follow provincial instructions for any required additional cleaning of the container prior to its disposal.
2. Make the empty container unsuitable for further use.
3. Dispose of the container in accordance with provincial requirements.
4. For information on the disposal of unused, unwanted product and the clean-up of spills contact the regional office of the Conservation and Protection Service, Environment Canada.

NOTICE TO USER: This control product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products Act to use this control product under unsafe conditions.

NOTICE TO BUYER: Seller's guarantee shall be limited to the terms set out on this label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

\*\*\*\*\*

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))