Avant Guard(R)
Flea Collar For Medium Dogs

with Integral Buckle

KILLS
FLEAS UP TO 4 MONTHS AND AMERICAN
DOG AND BROWN DOG TICKS UP TO 2 MONTHS

GUARANTEE:
Propoxur 9.4%

DOMESTIC

REG. NO. 23537 P.C.P. ACT

READ THE LABEL BEFORE USING
KEEP OUT OF REACH OF CHILDREN

Imported by:
Avant Sales
6991 Millcreek Drive, Unit 6,
Mississauga, On. L5N 6B9

30.2 g
Fits up to 52.5 cm Neck Size

Avant Guard(R)
Flea Collar
with Integral Buckle
For Medium Dogs

READ THE LABEL BEFORE USING.

DIRECTIONS: For best results, before placing this collar on,
bathe and shampoo pet with a pyrethrin based product such as
Avant Guard Dog Flea & Tick Shampoo. This will produce a

"quick kill" effect.

Remove the collar from the package and place around dog's neck. Thread flat end of the collar through buckle hole and tighten as necessary, so that at least two ridges are covered by the buckle, leaving ample room for expansion as the dog grows. Cut off any excess length and discard in garbage. Collar must be worn loosely, but securely enough to prevent easy removal. After collar placement, adult ticks will be killed in a few days and will fall off or may be easily removed. American Dog and Brown Dog ticks appear on dogs in three stages. In the first two stages, ticks are small and are difficult to see. The collar will also control these immature stages. Collar will kill ticks encountered in first 2 months of wear. Kills dog fleas for up to 4 months. FOR CONTINUOUS PROTECTION, REPLACE COLLAR WHEN EFFECTIVENESS DIMINISHES. To release the collar, squeeze between thumb and finger and pull sharply.

FLEAS: This collar is also effective in killing fleas. Temporary wetting will not reduce effectiveness. If your dog goes swimming or is out in the rain, it is not necessary to remove the collar. The continuous action collar will rapidly replace any flea and tick killing material removed. For effective flea and tick control, treatment of the pet should be combined with the sanitation of any area used by the pet. Vacuum floors, carpets, and furniture (discard vacuum bag after use) and wash the pet's bedding, living quarters, and surrounding areas. If pest problems persist, an insecticidal premise treatment may be required. For best results, use in combination with a complete flea eradication program.(See enclosed leaflet for more information.)

PRECAUTIONS: Do not open protective pouch until ready to use. DO NOT ALLOW CHILDREN TO HANDLE THIS COLLAR. Dust will form on the collar during storage. Do not get dust or collar in the mouth. Harmful if swallowed. Do not get dust in eyes. Wash hands thoroughly with soap and water after handling. Do not use on nursing dogs, or dogs under 8 weeks of age. When collar is first worn, observe neck area frequently for signs of irritation. Remove collar immediately if irritation or adverse reaction is noted. Collar is intended for use only as an insecticide generator and is not to be taken internally by humans or animals. Do not use on sick or convalescing dogs, or dogs under drug treatment. Do not use other cholinesterase inhibitors (other organophosphate or carbamate containing parasiticides) while collar is worn or for a few days after

removing collar. Before using on pregnant animals, consult a veterinarian.

FIRST AID: If dust gets in eyes or on skin, flush with plenty of water. If irritation develops or persists or if ingested, obtain medical attention or call Poison Control Centre.

TOXICOLOGICAL INFORMATION: Propoxur is a cholinesterase inhibitor. Atropine is antidotal.

DISPOSAL: Wrap and dispose of any excess collar, packaging and expired collars with garbage to keep out of reach of children.

****

+))