

86.12.19

Propaturf Herbicide

Liquid Emulsifiable Concentrate - For Turf

Net Contents 20 Litres

Contains: Dichlorprop and 2,4-D
low-volatile butoxy-ethanol esters

Commercial

Guarantee:

Dichlorprop present as butoxy-ethanol esters 350 g/L
2,4-D present as butoxy-ethanol esters 329 g/L

Registration No. 14800
Pest Control Products Act

Read the label before using

Warning Poison

May & Baker Canada Inc.
1274 Plains Road East
Burlington, Ontario
L7S 1V6

DIRECTIONS FOR USE

General Information: Propaturf is a specially formulated herbicide for golf course fairways and roughs, sod farms, grass areas in parks and parklands, and also in non-crop areas. Propaturf should not be used on Bentgrass.

Method of Application: The weeds listed below are most susceptible in May, June and September (in coastal areas of B.C.,

also in April and October). The effectiveness of a treatment in July and August depends on general weather conditions. If it is hot and dry during that period, a second treatment may be necessary. If it rains within 24 hours after spraying, a repeat application may be required. Newly seeded areas should not be sprayed within 4 weeks after grass emergence and then only at half the recommended rate for the first month.

Rate of Application: Use 3.5 litres of Propaturf in 200-500 litres of water per hectare. The amount of water needed to cover one hectare depends on the spray equipment and is not important, provided it is not less than 200 L/ha.

Weeds Controlled:

Black Medick	Plantain
Bull Thistle	Prostrate Knotweed
Burdock	Purslane
Canada Thistle	Ragweed
Chicory	Redroot Pigweed
Cinquefoil (topkill)	Round-leaved Mallow
Common Chickweed	Sheep Sorrel
Creeping Charlie	Shepherd's-Purse
Curled Dock	Sow-thistle
Dandelion	Speedwell
Goat's-beard	Stinkweed
Goldenrod	Tall Buttercup
Hawkweed	Toadflax
Horsetail	Teasel
Lady's-thumb	Vetch
Lamb's-quarters	White Clover
Mayweed	Wild Carrot
Milkweed (topkill)	Wild Strawberry
Mouse-eared Chickweed	Yarrow (may need 2nd treatment)
Mullein	Yellow Rocket
Mustard	
Oxalis	

Poison Ivy: can be controlled with Propaturf at the rate of 1 litre per 100 litres of water. Spray thoroughly until the foliage drips. The best results are obtained when applied between mid-June and late July. A repeat treatment in the following year may be necessary.

Precautions: Keep out of reach of children. Harmful if swallowed or inhaled. Avoid contact with skin, eyes or clothing. Sprays and vapours may injure broadleaf crops such as vegetables, fruits or ornamentals. Use only for recommended purposes and at recommended rates. Do not use the same spray equipment for other purposes. Keep in original container during storage. Do not contaminate water supplies, irrigation ditches, ponds, lakes or streams.

First Aid: This product contains petroleum distillates. If swallowed, do not induce vomiting. Rush patient to nearest hospital taking this labelled container with you. In case of skin contact, remove contaminated clothing and wash skin thoroughly with soap and water. If in eyes, flush with plenty of water for 15 minutes and get medical attention.

Storage and Disposal: May be stored at any temperature. Shake well after storing for one year or longer. To dispose of empty container, puncture and bury at least 50 cm deep and away from any water supplies.

Notice to User: This control product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products Act to use a control product under unsafe conditions.

Notice to Buyer: Seller's guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))