

31-DEC-1985

(R) BAYLUSCIDE
(FORMERLY BAY 73)

5% GRANULAR

SEA LAMPREY LARVICIDE
CONTAINS NICLOSAMIDE

FOR USE ONLY BY PERSONNEL OF
GOVERNMENT FISHERIES AGENCIES.

FOR USE IN SURVEY WORK TO DETERMINE
POPULATION LEVELS OF SEA LAMPREY LARVAE.

RESTRICTED

NET WEIGHT
KILOGRAMS

GUARANTEE:

NICLOSAMIDE.....5%

CANADIAN PATENT NO. 686, 211

REG.NO. 10,841
P.C.P. ACT

CAUTION
KEEP OUT OF REACH OF CHILDREN

(SEE REAR PANEL FOR FIRST AID AND CAUTION STATEMENTS)

READ THE LABEL BEFORE USING

CHEMAGRO LIMITED
1355 AEROWOOD DRIVE

MISSISSAUGA, ONTARIO,
CANADA
L4W 1C2

NOTICE TO USER: This control product is to be used only in accordance with the directions on this label. It is an offense under the PEST CONTROL PRODUCTS ACT to use a control product under unsafe conditions.

RESTRICTED USES

This product is to be used only in the manner authorized; contact local authorities regarding appropriate use permits that may be required.

DIRECTIONS FOR USE

BAYLUSCIDE 5% Granular has proved to be a useful tool for determining populations of Sea Lamprey larvae in streams. The granules, when applied to areas to be surveyed, fall rapidly to the stream bottom where they give excellent kill of lamprey eel larvae.

Apply 110 kg of BAYLUSCIDE 5% Granular evenly over each surface hectare of water (1.25 g/100 sq. m). Use equipment that can be accurately calibrated to distribute the required amount of granules evenly over the area to be treated.

When applied as directed, BAYLUSCIDE will not harm game fish populations that may be present in treated waters. BAYLUSCIDE applications will be toxic to freshwater clams and mussels and may be toxic to fish species that are physiologically bound to the stream bottom.

CAUTION

May be harmful if swallowed. Wash thoroughly with soap and water after handling. Do not contaminate feed or food. Keep out of reach of children.

FIRST AID: In case of poisoning, call a physician immediately. Have patient lie down and keep quiet. If swallowed, vomiting should be induced. Administer milk or water freely and induce vomiting by giving one dose(15 mL) of syrup of ipecac. If vomiting does not occur within 10 to 20 minutes, administer second dose. If syrup of ipecac is not available, induce vomiting by sticking finger down throat. Repeat until vomit fluid is clear. The patient should be lying down with the head below the foot level and facing down or to one side. Professional medical assistance should be secured immediately. DO NOT INDUCE VOMITING TO AN UNCONSCIOUS PERSON OR TO PERSONS IN A CONVULSIVE STATE. If on skin, remove contaminated clothing and wash skin immediately with soap and warm water. If eyes are contaminated, wash immediately with flowing water for at least 15 minutes.

TOXICOLOGICAL INFORMATION: The exact toxic mode of action of BAYLUSCIDE is not known. In experimental animals acutely poisoned with BAYLUSCIDE no characteristic symptoms were noted. The patient should be treated symptomatically.

NOTICE

Seller's guarantee shall be limited to the terms set out on the label and, subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

BAYLUSCIDE IS A REG.TM OF BAYER AG.

Printed in U.S.A.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this

service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))