

85.12.31

ORTHO

FLOTOX*

Garden Sulphur W.P.

For Disease and Insect Control on Fruit Trees, Grapes
and Ornamentals. (Dust on Ornamentals & Tomatoes)

READ THE LABEL BEFORE USING.

GUARANTEE

SULPHUR 90%

REG. NO. 14090

P.C.P. ACT

*Trademark of Chevron Chemical Co.

DOMESTIC

NET CONTENTS 709g

READ THE LABEL BEFORE USING

USE STRICTLY IN ACCORDANCE WITH LABEL
CAUTIONS & DIRECTIONS

Chevron Chemical (Canada) Limited
3228 South Service Road
Burlington, Ontario
L7N 3H8

KEEP PESTICIDE IN ORIGINAL CONTAINER. DO NOT PUT CONCENTRATE OR
DILUTE INTO FOOD OR DRINK CONTAINERS.

Plants	Pests	Comments
--------	-------	----------

ROSES	Powdery Mildew, Black Spot Rust. Combination spray to control Powdery Mildew, Black Spot Rust, Aphids, Red Spiders, Thrips and Leafhopper.	Dust or spray every 7-10 days during the growing season 50g (3 Tablespoonfuls), FLOTOX Garden Sulphur 15 mL (1 Tablespoonful) ISOTOX Insect Spray per 5L of water.
DAHLIAS, CHRYSANTHEMUMS PEAS, BEANS	Powdery Mildew, Black Spot Rust	Dust or spray every 7-10 days during growing season. Preharvest interval for Peas and Beans is 1 day. Apply when infection appears.
GRAPES	Powdery Mildew	Spray when mildew first appears and repeat after 10 days. Continue spray as necessary. Preharvest interval 1 day except when grapes are used for wine then the interval is 21 days. Do not use on sulphur-sensitive varieties such as Concord.
SWEET CHERRIES	Brown Rot, Leaf Spot, Powdery Mildew	For Brown Rot and Leaf Spot spray FLOTOX from bloom up to 1 day before harvest. If weather is wet at harvest spray between pickings. For Powdery Mildew spray at husk-fall and repeat in 10 days.

SOUR CHERRIES	Brown Rot, Blossom Blight, Powdery Mildew, Rust Mite	Apply FLOTOX at bloom for Blossom Blight. For Powdery Mildew control spray at first, second and third cover. For Rust Mites apply FLOTOX in summer sprays. Preharvest interval 1 day.
PEACHES, PLUMS	Brown Rot, Leaf Spot Scab, Powdery Mildew (Peach), Brown Rust (Plum), Black Knot (Plum)	For Brown Rot, Leaf Spot, Brown Rust, Black Knot and Scab control apply spray at pink, bloom, shuck, cover and pre/harvest. If wet during picking, spray between pickings. For Powdery Mildew control in Peaches, spray at husk-fall and again in 2 weeks. FLOTOX can be used up to harvest.

<p>APPLES, PEARS</p>	<p>Powdery Mildew, Scab, Rust Mites</p>	<p>Apply a spray in a regular protective schedule from green tip up to and including first cover spray. Depending on mildew conditions and temperature, rates may be slightly reduced after the first cover. During wet weather additional treatments may be needed. Apply FLOTOX in summer sprays for Rust Mite control. Preharvest interval 1 day.</p>
<p>TOMATOES</p>	<p>Russet Mite</p>	<p>Dust or spray when fruit begins to set. Repeat treatments as necessary. Preharvest interval 1 day.</p>
<p>CURRANTS</p>	<p>Powdery Mildew, Rust</p>	<p>Spray immediately before the earliest blossoms open and continue applications through harvest. Preharvest interval 1 day.</p>
<p>GOOSEBERRIES</p>	<p>Powdery Mildew</p>	<p>Spray immediately before the earliest blossoms open and continue applications through calyx. Do not use on sulphur sensitive varieties.</p>

As a Spray-Mix 50g (3 Tablespoonfuls) FLOTOX Garden Sulphur in a small amount of water, stirring until a thin paste is formed, then add water to make 5 litres of mixture. Stir thoroughly. Do not store diluted spray. Make new dilution for each use.

FIRST AID: Avoid contact with skin or eyes. In case of eye contact, immediately flush eyes with plenty of water. Get medical attention if irritation persists. Wash hands after use. If swallowed, induce vomiting. Repeat until vomit is clear. Obtain medical attention.

PRECAUTIONS: Do not use within 4 weeks of an oil spray treatment. Sulphur may cause injury to foliage if applied when temperature is exceedingly high. Do not make applications at this time. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not reuse container. Destroy when empty. KEEP OUT OF REACH OF CHILDREN.

NOTICE TO USER: This control product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products Act to use a control product under unsafe conditions.

LIMITATION OF WARRANTY: Sellers guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to person or property arising from the use or handling of this product and accepts the product on that condition.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))