

2000/2002

**ESTAKIL MCPA 500  
LIQUID HERBICIDE**

Contains MCPA Emulsifiable Concentrate

FOR CONTROL OF BROADLEAF WEEDS  
IN BARLEY, WHEAT, RYE AND FLAX.

AGRICULTURAL

READ THE LABEL BEFORE USING

GUARANTEE:

MCPA (present as mixed butyl ester) ...500 grams per litre

REGISTRATION NO. 14662 PEST CONTROL PRODUCTS ACT

**WARNING      POISON**

NET CONTENTS: 10 LITRES

Agrolinz, Inc.  
c/o Nufarm, Inc.  
500 Lower Lake Road  
St. Joseph, MO 64504 USA

Canadian Agent  
Nufarm Agriculture Inc.  
5507 1<sup>st</sup> Street SE  
Calgary, Alberta T2H 1H9

24 hours emergency Response Number 1-202-483-7616

**DIRECTIONS FOR USE**

This product may be applied by aerial or conventional ground spraying equipment. Aerial spraying is a specialized job, hence only competent, reliable operators should be employed. Apply lower rates to weed seedlings in 2 to 4 leaf stage which are growing rapidly and higher rates for more advanced weeds or during dry and cool weather or heavy infestation.

| CROP                     | RATE PER HECTARE | HA TREATED BY 10 L |
|--------------------------|------------------|--------------------|
| Barley, Oats, Rye, Wheat | 1.25-2.25 L      | 4.5-8              |

DIRECTIONS: Apply sufficient water to obtain good coverage. Treat spring seeded cereals when crop is in 2-4 leaf stage, combining maximum crop tolerance and weed susceptibility. May be applied at early shot blade stage and again from the milk stage to full maturity. For fall seeded wheat or rye, treat when crop growth begins until early shot blade stage. Use lower rates for cereals under-seeded to legumes.

| CROP | RATE PER HECTARE | HA TREATED BY 10 L |
|------|------------------|--------------------|
| Flax | 1.25-2.25 L      | 4.5-8              |

DIRECTIONS: Spray crop when plants are 5 cm high till pre-bud stage.

| CROP  | RATE PER HECTARE | HA TREATED BY 10 L |
|---|------------------|--------------------|
| Established Alfalfa, Alsike, Red Clover and grasses | 700 mL-2.0 L     | 5-14.5             |

DIRECTIONS: Apply 150 to 200 liters of water per hectare. Treat before the development of legumes has occurred and when they are covered by a canopy of weeds and crop. Do not apply in the presence of sweet clover.

| CROP   | RATE PER HECTARE       | HA TREATED BY 10 L |
|--|------------------------|--------------------|
| Uncropped Land (drainage ditches, pastures, roadsides) | 1.75-3.25 L<br>2.2-4.5 | 3-6<br>2.25-4.5    |

DIRECTIONS: For general weed control, apply 100-200 liters of water per hectare. Treat in early spring or late fall when weeds are in bud. The low rates may be used on more susceptible weeds. For control of woody plants (poplars, western snowberry, and willows), treat when plants are in full leaf.

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN. DO NOT SPRAY during windy conditions or when air temperature is above 27C as vapor or spray mist may drift on desirable broadleaf plants. Susceptible plants include beans, sugar beets, tomatoes, rape, sunflowers, peas, clovers trees and shrubs. Hazard of drift can be reduced by coarse sprays, low spraying pressure. DO NOT CONTAMINATE

domestic or irrigation waters. Thoroughly clean sprayer after use. DO NOT STORE near or in the same room as seed, fertilizers, insecticides or fungicides. Allow 7 days between application of product and grazing/cutting for hay.

FIRST AID: Avoid contact with skin, eyes and clothing. If swallowed, do not induce vomiting, but rush patient to nearest hospital taking this container or contact a Poison Control Centre. This product contains petroleum distillates. In case of contact on skin, wash with plenty of soap and water, if in eyes, flush with plenty of water and get medical attention or contact a Poison Control Centre.

STORAGE: May be stored at any temperature. Shake well after storing for one year or longer.

DISPOSAL:

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple - or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER: This control product is to be used only in accordance with the directions on this label. It is an offense under the Pest Control Products Act to use a control product under unsafe conditions.

NOTICE TO BUYER: Seller's guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

\*\*\*\*\*

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))