

85.12.31

weed
away

2,4-D Amine
500

LIQUID HERBICIDE

AGRICULTURAL

READ THE LABEL BEFORE USING

GUARANTEE: 2,4-D 500 g/L
(PRESENT AS AMINE)

REGISTRATION NO. 17411
PEST CONTROL PRODUCTS ACT

20 LITRES

WARNING POISON

INTERPROVINCIAL COOPERATIVE LIMITED. P.O. BOX 1050,
SASKATOON, SASKATCHEWAN S7K 3M9

WINNIPEG TORONTO MONTREAL MONCTON

DIRECTIONS:

For ground or air application - Add water to spray tank until 1/2 full. Add 2,4-D Amine 500, then complete filling with water and mix thoroughly. Treat weeds when small for best results and least cost. Use 60-240 litres of water per hectare unless otherwise specified.

WHEAT, BARLEY, RYE:

Treat from third leaf to early shot blade stage. Do not treat cereals underseeded to legumes.

EASY-TO-KILL WEEDS: Mustard, Lambs Quarters, Shepherd's Purse, Stinkweed, Flixweed, Ragweed, Blue Bur, Kochia, Prostrate Pigweed, Cockle Bur, Prickly Lettuce. Small Seedlings growing rapidly: 700 mL/ha. Weeds in bud, dry or cool weather. Heavy infestation: 1.1 L/ha.

HARDER-TO-KILL WEEDS: Red Root Pigweed, *Sow Thistle, *Canada Thistle, *Leafy Spurge, *Blue Lettuce, Dog Mustard, Common Peppergrass, Russian Thistle. (*) Top growth suppression only. Small seedling growing rapidly: 1.25 L/ha. Weeds in bud, dry or cool weather. Heavy infestation: 1.75 L/ha.

OATS

(Emergency use only). Treat from emergence to two leaf stage or from the six leaf to early shot blade stage. 2,4-D reduces oat yields. Use MCPA unless weeds resistant to MCPA are present. 2,4-D Amine is safer than 2,4-D Ester. Dosages above 1 L/ha. may cause injury.

FLAX

Linseed (Emergency use only). Treat from three leaf stage to before budding. MCPA is safer to use than 2,4-D, especially above 700 mL/ha. Dosages above 1 L/ha. may cause injury.

FALL WHEAT:

Treat in Spring only, from full tillering to shot blade stage. Do not treat wheat underseeded to legumes. Dosages above 1.1 L/ha. may cause injury. SUSCEPTIBLE WEEDS: Small growing seedlings: 700 mL/ha. Bud stage, adverse weather: 1 L/ha. RESISTANT WEEDS: Small growing seedlings: 1.1 L/ha. Bud stage, adverse weather, : 1.1 L/ha.

GRASS PASTURES: (ESTABLISHED)

Treat in Spring or Fall for best results. Use 50-400 litres of water per hectare. Susceptible weeds: 2 L/ha. Intermediate susceptibility (adverse conditions): 3.5 L/ha.

CORN:

Treat before corn is 15 cm tall. Controls Atrazine resistant Lambs Quarters and other susceptible broadleaf weeds. Use 100-200 litres of water per hectare. Weeds, small growing actively: 700 mL/ha. Weeds, larger adverse weather: 1 L/ha.

WOODY PLANTS:

Foliage application: 4.25-8.5 L in 1000 L of water per hectare.

TANK MIXES WITH 2,4-D AMINE 500

Use tank mixes only when they are registered or where recommended. Read and follow directions on both products used in a tank mix.

1. 2,4-D Amine 500 at 600-800 mL per hectare (300-400 g active per hectare) plus Dicamba, 150 grams active per hectare on wheat or 100 grams active per hectare on barley. Weeds controlled include Canada and Sow Thistle, Buckwheat, Smartweed and Cow Cockle.

2. 2,4-D Amine 500 - use up to 1.1 Litres per hectare (550 g active) in combination with TCA at 1.1 kilograms active per hectare. Use only on barley. Apply at the 4 leaf stage of the crop when weeds are in the 1-3 leaf stage. Use 100-150 litres of water per hectare. Add and dissolve TCA before 2,4-D is added. Weeds controlled - 2,4-D susceptible broadleaf weeds plus Green Foxtail.

PRECAUTIONS:

KEEP OUT OF REACH OF CHILDREN. Avoid contact with skin, eyes or clothing. Remove and wash contaminated clothing and shoes before re-use. Do not contaminate streams, lakes, ponds, irrigation ditches or waters used for livestock or domestic purposes. Keep in original container during storage. Do not store near seeds, fertilizers, insecticides or fungicides. Do not store at temperatures below freezing. Do not spray on windy days or allow spray, mist or vapours to contact plants other than those being treated. Do not use sprayers which have contained 2,4-D to apply other pesticides to crops.

FIRST AID:

On skin, wash immediately with soap and water. In eyes, flush with water for 15 minutes, then seek medical attention.

DISPOSAL PROCEDURES

Destroy empty container. Dispose of wastes by burying in a non-crop area away from all water supplies. Follow local regulations if they have been established.

NOTICE TO USER:

This control product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products Act to use a control product under unsafe conditions.

NON-WARRANTY

Seller's guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not

liable for any loss resulting, directly or indirectly, from
reliance upon this service.

+))