

11-APR-2007

18-DEC-2007 Notification - Change in Registrant Address

14-FEB-2008 Notification - Change in Registrant Name

TOTALEX®

GRASS AND WEED KILLER READY -TO- USE

Brush, Grass & Broadleaf Weed Control Liquid
Controls Entire Plant, Burns Off Leaves And Stem And Kills Roots

DOMESTIC
KEEP OUT OF REACH OF CHILDREN
NON-SELECTIVE HERBICIDE

GUARANTEE: Glyphosate 7 g/L (present as isopropylamine salt)

Contains 1,2 benzisothiazolin-3-one at 0.04% as a preservative.

REGISTRATION NO. 26610.01 PEST CONTROL PRODUCTS ACT

NO RESIDUAL ACTIVITY IN SOIL

USES: ALONG FENCE, EDGING/TRIMMING LAWNS, PATIO WEED CONTROL, DRIVEWAYS,
LAWN/GARDEN RENOVATION, HARD-TO-MOW AREAS (SEE DIRECTIONS).

READ THE LABEL BEFORE USING

Teragro Inc.

Notification Change

NET CONTENTS 1 L

~~Virterra Products Corporation~~

~~R.R. 7, Site #7, Box 16~~

~~Calgary, Alberta T2P 2G7~~

~~(403) 285-0500 www.virterraproducts.com~~

P.O. Box 137

Chestermere AB T1X 1K8

**In case of a medical emergency involving this product, call toll free day or night
1-800-228-5635, x135**

® GLYFOS is a registered trademark of Weed-Master Inc.

Not for reformulation or repackaging.

PRECAUTIONS FOR USE

Avoid spray contact with desirable plants as any plant contacted may be killed. Do not spray in windy conditions. Do not spray to the point of run-off. Heavy rainfall immediately after application may wash the chemical off the foliage and repeat treatment may be required. Do not apply if rainfall is forecast for the time of application.

PRECAUTIONS

CAUSES EYE IRRITATION. HARMFUL IF SWALLOWED. Avoid contact with eyes or prolonged contact with skin. Do not allow others such as children and pets on treated areas during application or to re-enter treated areas until spray has dried. Wash immediately after using this product. Do not contaminate foodstuffs, seed or feed by storage or disposal.

FIRST AID

If on skin, flush with water.

If ingested, drink water or milk and get medical attention or call a Poison Control Centre.

If in eyes, immediately flush with plenty of water for at least 15 minutes. Call a physician or contact a Poison Control Centre.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION: Treat symptomatically.

DISPOSAL

Do not reuse empty container. Dispose of empty container with household garbage.

GENERAL PRODUCT INFORMATION

TOTALEX GRASS AND WEED KILLER READY TO USE is a non-selective herbicide. It controls most annual and perennial grasses, including lawn grasses, broadleaf weeds and brush (i.e. virtually anything that is green and growing). It is absorbed by the leaves and moves throughout the stem and roots to control the entire plant. Mature perennial weeds should be treated after seed heads, flowers or fruit appear. All plants are most easily controlled in the young, actively growing, seedling stage.

TOTALEX GRASS AND WEED KILLER READY TO USE wilts and browns plants within 48 hours. Plants usually dry up within 7 to 14 days. TOTALEX GRASS AND WEED KILLER READY TO USE has no soil activity and will not run off to affect nearby vegetation. To ensure that TOTALEX GRASS AND WEED KILLER READY TO USE gets down to the roots, avoid tillage or disturbing the treated area for 7-10 days after treatment or as directed below. Make sure to contact only those plants that you want to kill. TOTALEX GRASS AND WEED KILLER READY TO USE will damage or kill any desirable plants it contacts. Use a repeat application on any seedlings that re-grow from seeds.

DIRECTIONS

USE	APPLICATION INSTRUCTIONS	CONTROLS	MIXING/ COVERAGE
Undesirable vegetation control around buildings, lawn trimming/edging, patio weed control, vacant lots, storage & recreational areas, driveways, hard to mow areas and other similar areas	Apply to vigorously growing plants. Apply to annual weeds anytime after emergence. For perennial weeds apply after seed heads, flowers or berries appear.	Most annual and perennial grasses and weeds, such as quackgrass, chickweed, ragweed, knotweed, poison ivy, Canada thistle, milkweed, bindweed.	Ensure uniform and complete spray coverage but not to the point of runoff. This 1 L container treats a total of 20 m ² (5 paces X 4 paces*) * 1 pace = approx. 1 m
Around trees/shrubs/ ornamentals	Apply around base in a neat fashion. Avoid spray or drift contact with foliage, stems or green bark of desirable plants.		
Lawn/garden renovation	For best results on lawns, skip one mowing prior to treatment. Allow at least 3 days after treatment before disturbing the soil and/or replanting.		

Brush control	Apply after leaves are fully expanded. Treat trees and brush up to 1.5 m in height only; for taller trees and brush, cut back to 1.5 m or cut down and spray regrowth.	Most brush such as poplar, alder, maple and raspberry	
---------------	--	---	--

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.