dispar

DISBLOK

BAIT BLOCK

GUARANTEE: CHLOROPHACINONE   0.005%

READ THE LABEL BEFORE USING

DIRECTIONS FOR USE

FOR NORWAY AND ROOF RATS PLACE BAITS DISBLOK AREAS VISITED BY
RODENTS SUCH AS: TUNNELS, PATHS AND WHERE THEY USUALLY EAT.
REPLACE ANY STALE OR MOULDY BAITS AND BAIT FOR 15 DAYS AND
CONTINUE BAITING UNTIL ALL SIGNS OF FEEDINGS HAVE STOPPED. WHERE
A CONTINUOUS SOURCE OF INFESTATION IS PRESENT, PERMANENT DISBLOK
STATIONS SHOULD BE ESTABLISHED AND THE BAIT DISBLOK REPLACED AS
NEEDED

PRECAUTIONS: KEEP BAIT OUT OF REACH OF CHILDREN, PETS AND
LIVESTOCK.

REGISTRATION NO. 19036 PEST CONTROL PRODUCTS ACT

DISPAR VETERINARY PRODUCTS
JOLIETTE, QUEBEC,
CANADA
J6E 3Z1

NOTICE TO USER:

This control product is to be used only in accordance with the
directions on this label. It is an offence under the Pest
Control Products Act to use a control product under unsafe
conditions.

PRECAUTIONS:

Keep out of reach of children. If taken in large doses by humans
or animals, it may reduce the clotting ability of the blood and
cause internal bleeding. Place baits in protected places
inaccessible to children and domestic animals or in tamper-proof
bait boxes. Do not contaminate food, feed and drinking water.
Burn or bury bodies of dead rodents.

FIRST AID INSTRUCTIONS:

If swallowed give 1 or 2 glasses of water to patient and induce
vomiting placing finger in throat. Repeat until vomit is clear.
Call physician.

TOXICOLOGICAL INFORMATION:

Blood transfusions combine with intravenous injections and oral
doses of vitamin K1 are indicated as if it was haemorrhage caused
by an excessive dose of dicumarol.

****

+))