

97.01.30

NET CONTENTS
22.68 KG

MICRO/NIASUL W

FUNGICIDE
SULPHUR WETTABLE POWDER

AGRICULTURAL

GUARANTEE: Sulphur 92%

REGISTRATION NO. 14713
PEST CONTROL PRODUCTS ACT

READ THE LABEL BEFORE USING

WILBUR-ELLIS Company
P.O. Box 16458
Fresno, CA
93755

DIRECTIONS FOR USE

GENERAL INFORMATION: Apply in sufficient water for thorough coverage of all plant parts; this volume depends on the spray equipment and the size of trees and vines. Follow Provincial recommendations for proper timing of the sprays. DO NOT APPLY MICRO-NIASUL if high temperatures (above 27°C) and high humidity prevail or are expected during the next three days following application.

CROP/DISEASE	METHOD AND RATE
APPLE, PEAR Powdery Mildew Scab	Apply 6.5 kg/1000 L or 21.5 kg/ha in the Pre-Bloom, Calyx and First Cover sprays. In the later Cover sprays use 3.5-4.5 kg/1000 L or 11.5-15 kg/ha depending on mildew conditions and temperature.

PEACH Brown Rot Scab	Apply 6.5 kg/1000 L or 21.5 kg/ha in a regular spray program starting at the late Pink stage. Repeat just before and during harvest if wet weather occurs.
PLUM Brown Rot Black Knot	Apply 6.5 kg/1000 L or 21.5 kg/ha in a regular spray program starting at the Schuck stage. Repeat just before and during harvest if wet weather occurs.
RUTABAGAS Powdery Mildew	Apply 6.8 kg/ha in 675 to 1125 L of water/ha depending on the size of the rutabagas. Thorough coverage of both upper and lower leaf surfaces is essential for powdery mildew control. Apply weekly, starting when disease symptoms first appear in the field. Use no more than (5) applications per season.
SWEET CHERRY Brown Rot	Apply 6.5 kg/1000 L or 21.5 kg/ha in a regular spray program from the Bloom stage until just before harvest. Repeat during harvest if wet weather occurs.
SOUR CHERRY Brown Rot Powdery Mildew	Apply 6.5 kg/1000 L or 21.5 kg/ha at the Bloom stage. If Powdery Mildew is problem, apply 3.5 kg/1000 L or 11.5 kg/ha in the First, Second and Third cover sprays.
GRAPE Powdery Mildew	Apply 4.5 kg/1000 L in the Post-Bloom spray on Agawam, Seneca and Seibel varieties. Do not use on Concord, Foch and Van Buren varieties. Apply 6 kg/1000L in the First, Second and further Cover sprays. Do not apply later than 21 days before harvest on grapes for wine.
GREEN HOUSE CUCUMBER Powdery Mildew	Apply 1.5 g/L to run-off using a backpack of other suitable sprayer, at weakly intervals as needed to control Powdery Mildew. Leave at least 1 day between last spray and harvest.
PEAS Powdery Mildew	Apply 1.5 Kg/ha in 150 L of water. Apply when conditions are suitable for disease development or disease has appeared. Repeat weekly. Provide thorough coverage of upper and lower leaf surfaces. Do not apply more than 5 times per season. Leave at least 1 day between last spray and harvest.

COMPATIBILITY: This product is compatible with most commonly used insecticides and fungicides, except DNOC (Elgetol) and oil.

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN. Do not inhale dust of spray mist. Avoid contact with skin and eyes. After handling, wash thoroughly with soap and water before eating or smoking. Do not contaminate ponds, streams or other bodies of water. Avoid contamination of feed or food.

FIRST AID: If swallowed, induce vomiting by touching the back of the throat with a finger or blunt object; repeat until vomit fluid is clear. Take patient to nearest hospital taking this labelled container with you. In case of skin contact, remove contaminated clothing and wash skin with soap and water. If in eyes, flush with plenty of water for 15 minutes and go to nearest hospital.

STORAGE: Store in a cool, dry place away from food and feed.

DISPOSAL:

- (1) Empty container (bag) completely.
- (2) Follow provincial instructions for any required additional cleaning of the container prior to its disposal.
- (3) Make the empty container unsuitable for further use.
- (4) Dispose of the container in accordance with provincial requirements.
- (5) For information on the disposal of unused, unwanted product and the cleanup of spills, contact the Provincial Regulatory Agency or the Manufacturer.

NOTICE TO USER: This control product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products Act to use a control product under unsafe condition.

NOTICE TO BUYER: Seller's guarantee shall be limited to the terms set out on the label and subject thereto the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))