

93.06.22

((BOX))

Ralston Country Roads (R)

FLEA & TICK COLLAR FOR SMALL DOGS

Integral Buckle
(No Metal)

REGISTRATION NO. 21333 PEST CONTROL PRODUCTS ACT

DOMESTIC

Kills fleas up to 4 months

READ LABEL INSERT BEFORE USING

GUARANTEE: Propoxur 9.4%

NET CONTENT: 32.7g

SANDOZ AGRO CANADA INC.
Suite 302, Plaza 4, 2000 Argentia Road
Mississauga, Ontario L5N 1W1

Manufactured for:

Ralston Purina Canada Inc.
Woodstock, ON, N4S 7X5

PRECAUTION: Do not open protective pouch until ready to use. DO NOT ALLOW CHILDREN TO HANDLE THIS COLLAR. Dust will form on this collar during storage. Do not get dust or collar in mouth, harmful if swallowed. Do not get dust in eyes. Wash hands thoroughly with soap and water after handling.

Do not use on dogs younger than 3 months old or on nursing females. Consult your veterinarian before using this product on pregnant animals.

When collar is first worn observe neck area every few days for irritation. Remove collar at the first sign of irritation or adverse reaction. Collar is intended for use only as an insecticide generator and is not to be taken internally by man or animals. Do not use on sick or convalescing dogs. Do not use other pesticides on dog or administer other cholinesterase inhibitors (e.g. oral parasiticides), while collar is worn or for a few days after removing collar. Dispose of any excess collar, packaging and expired collars with garbage to keep out of reach of children.

FIRST AID: In case of poisoning, call a physician immediately. If dust from collar contacts skin, wash thoroughly with soap and water. If dust contacts eyes, flush eyes with water and get medical attention.

TOXICOLOGICAL INFORMATION: Dust in eyes will cause temporary pupillary constriction. Dust release from collar is a cholinesterase inhibitor.

Atropine and homatropine are antidotal.

((INSERT))

Ralston country Roads (R)

FLEA & TICK COLLAR FOR SMALL DOGS

Integral Buckle
(No Metal)

Kills fleas up to 4 months

DOMESTIC

REGISTRATION NO. 21333 PEST CONTROL PRODUCTS ACT

GUARANTEE: Propoxur 9.4%

SANDOZ AGRO CANADA INC.
Suite 302, Plaza 4, 2000 Argentia Road
Mississauga, Ontario L5N 1W1

Manufactured for:

Ralston Purina Canada Inc.
Woodstock, ON, N4S 7X5

For use in the control of cat and dog fleas and American dog and American brown ticks on dogs.

DIRECTIONS: Remove the collar from the package and place around dog's neck. Thread flat end of collar through buckle hole and tighten as necessary, so that at least two ridges are covered by the buckle. Leaving ample room for expansion as the dog grows, cut off any excessive length and discard in garbage. Collar must be worn loosely, but securely enough to prevent easy removal. To release the collar, squeeze buckle between thumb and finger and pull sharply. Collar should be removed when dog is bathed with soap or detergent.

The collar starts killing fleas and ticks when it is placed around the dog's neck. The fleas and ticks on the dog will be killed and new ones which may temporarily appear on the dog, will also be killed during the 4 months (17 weeks) collar is worn. For continuous protection, replace collar when effectiveness diminishes.

PRECAUTION: Do not open protective pouch until ready to use. DO NOT ALLOW CHILDREN TO HANDLE THIS COLLAR. Dust will form on this collar during storage. Do not get dust or collar in mouth, harmful if swallowed. Do not get dust in eyes. Wash hands thoroughly with soap and water after handling.

Do not use on dogs younger than 3 months old or on nursing females. Consult your veterinarian before using this product on pregnant animals.

When collar is first worn observe neck area every few days for irritation. Remove collar at the first sign of irritation or adverse reaction. Collar is intended for use only as an insecticide generator and is not to be taken internally by man or animals. Do not use on sick or convalescing dogs. Do not use other pesticides on dog or administer other cholinesterase inhibitors (e.g. oral parasiticides), while collar is worn or for a few days after removing collar. Dispose of any excess collar,

packaging and expired collars with garbage to keep out of reach of children.

FIRST AID: In case of poisoning, call a physician immediately. If dust from collar contacts skin, wash thoroughly with soap and water. If dust contacts eyes, flush eyes with water and get medical attention.

TOXICOLOGICAL INFORMATION: Dust in eyes will cause temporary pupillary constriction. Dust release from collar is a cholinesterase inhibitor.

Atropine and homatropine are antidotal.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))