

1998/2003

SC Johnson (R)

Raid (R)

FLEA KILLER

PLUS

DOG SPRAY

DOMESTIC

GUARANTEE:

(S)-Methoprene	0.25%
Pyrethrins	0.18%
Piperonyl Butoxide, Technical	0.36%
N-Octylbicycloheptene Dicarboximide	0.60%

REG. NO. 23285

P.C.P. ACT

KEEP OUT OF REACH OF CHILDREN

READ THE LABEL BEFORE USING

WARNING EYE IRRITANT

CAUTION FLAMMABLE

175 mL

GENERAL INFORMATION: Kills adult dog fleas, American dog and brown dog ticks. Effectively breaks the flea life cycle and prevents the growth of the pre-adult stages of the flea. One treatment kills adult fleas for up to 7 days and keeps flea eggs from hatching for 30 days. Aids in control of flea bite dermatitis by killing fleas.

DIRECTIONS FOR USE: SHAKE WELL BEFORE USING. Cover animal's eyes with hand and with a firm fast stroke, spray head, ears and chest until damp but no soaked. With fingertips, rub into face around

mouth, nose and eyes. Then spray the neck, middle and hind quarters finishing with legs. Make sure spray thoroughly wets ticks. For best results, spray against natural lay of the hair, rub in well and, following the natural lay of the hair, wipe off excess with dry cloth and then brush. Using this technique, the application of 4 to 7 mL of Raid Flea Killer Plus Dog Spray per kilogram of body weight will provide effective flea control. Repeat only as necessary. Given the residual control afforded by Raid Flea Killer Plus, the application of this product once per month should be adequate. In exceptional circumstances, where there is a heavy external source of adult fleas re-infesting the animal, re-treatment may be increased to once per week.

PRECAUTIONARY STATEMENTS: KEEP OUT OF REACH OF CHILDREN. Harmful if swallowed. Avoid contact with eyes. After using, wash hands with soap and water. Do not treat puppies less than 8 weeks old or nursing females. Consult veterinarian before using on pregnant animals. Do not use on animals under drug or pesticide treatment. Do not use on sick animals.

FIRST AID: IF SWALLOWED, call physician immediately or Poison Control Centre. Drink 1 to 2 glasses of water and induce vomiting by touching the back of the throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. IF IN EYES, flush with plenty of water for 15 minutes. Get medical attention if irritation persists. IF ON SKIN, wash affected area with soap and water.

STORAGE AND DISPOSAL: Flammable. Keep away from heat or open flame. Do not reuse empty container. Wrap empty container and place in garbage.

Questions? Comments? Call 800-558-5566 weekdays 9-9 Eastern Time or write Helen Johnson.

(R) Trade Mark Reg'd

S.C. JOHNSON & SON, INC.
Reg'd User S.C. JOHNSON & SON, LIMITED.
Brantford (ON)
Canada N3T 5R1

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label

information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))