C-I-L®

## Cutworm and Maggot Killer - Granular

## **LARVEX**®

DOMESTIC

READ THE LABEL BEFORE USING KEEP OUT OF REACH OF CHILDREN

CAUTION

POISON

REG. NO. 13517 P.C.P. ACT

GUARANTEE: Diazinon 5%

250 g

Nu-Gro IP Inc. 10 Craig St., Brantford ON N3R 7J1

C-I-L® Cutworm and Maggot Killer (diazinon) is an effective control for cutworms, maggots, chinch bugs and other soil insects on lawns, vegetable gardens and around dwellings.

**DIRECTIONS:** 4 g = 1 teaspoon 12 g = 1 tablespoon

**VEGETABLE GARDENS:** For control of root maggots and other pests of vegetables, follow the directions given below. To be effective, the treatment must be uniformly applied. Some injury may occur to crops grown on sandy soils following the seed furrow treatments. The danger of such injury can be lessened if about 5 mm of soil separates the seed from the insecticide. Too much soil may decrease the effectiveness of the treatment. This

product will not be effective for controlling root maggots that have developed a resistance to diazinon.

Carrots and Parsnips - Carrot Rust Fly Larvae: Apply 10 q per 10 m row length in seed furrow at time of planting. Onions (including Green Bunching Onions) - Onion Maggots: Apply 10 q per 10 m row length in seed furrow at time of planting. Radishes - Root Maggots: Apply 18 g per 10 m row length in seed furrow at planting time. Two weeks after thinning, make a second application in a narrow band directly over the row. Root Maggots: Apply 43 g per 10 m row length (follow directions given for radishes). Cabbage, Cauliflower, Broccoli, Brussels Sprouts - In Seed Beds - Root Maggots: Apply 15 g per 10 m row length in seed furrow at planting time. As Transplants - Root Maggots: Apply 2 g per plant in a 10 cm radius around the stem of each plant soon after transplanting. Corn-Corn Rootworm: Apply 20 g in a 20 cm band over a 10 m row length at planting Tomatoes - Vinegar Flies: Broadcast 8 g per 10 m<sup>2</sup> of soil surface of the tomato patch when flies first appear. Repeat at 7-10 day intervals or as required. **Vegetables - Cutworms (surface** and subterranean): Uniformly distribute 50 g per 10 m<sup>2</sup> of soil surface and then mix into the top 8 to 10 cm of soil for surface cutworms and 8 to 15 cm for subterranean cutworms.

## LAWNS: FOR SPOT TREATMENT ONLY.

For the control of chinch bugs - Apply 20 g per m² in areas of infestation only. Water lightly before and after treatment.

Earwigs - Apply 20 g per m² in areas of infestation only. Lawn moth (sod webworm), sowbugs, and springtails (collembola). Apply 7.5 to 15 g per m² in areas of infestation only. Ants - Apply 3 g of granules over and around each individual ant hill. Digger wasps - Apply 3 g over and around each individual ground nest opening. Make application about dusk, when the wasps are not active.

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN. May be fatal if swallowed. Avoid inhaling dust or contact with eyes, skin and clothing. When applying to lawns wear a long-sleeved shirt, long pants, shoes and socks and chemical resistant (e.g. rubber) gloves during handling, application and watering-in. Rinse gloves before removal. Do not treat areas greater than 450 m² (5000 sq. ft.) In one day. Wash thoroughly after use. Avoid contamination of food, feed, utensils, drinking water or any body of water. Use only as recommended on label. This product is toxic to fish and wildlife. Keep people off treated lawns for 24 hours following application and watering in.

FIRST AID: If swallowed, induce vomiting. Contact a poison

control centre and obtain prompt medical attention. **If in eyes,** flush with water for 15 minutes and get medical attention. **If on skin,** wash with soap and water. Take container, label or product name and P.C.P. Registration Number with you when seeking medical attention.

**TOXICOLOGICAL INFORMATION:** Stomach lavage. Diazinon is a cholinesterase inhibitor. Antidote - Atropine sulphate at 1 mg to 2 mg every 15 minutes may be necessary.

DISPOSAL: DO NOT REUSE EMPTY CONTAINER. Dispose of empty container with household garbage. For additional product information or in case of emergency, spills or fire, call toll-free 1-800-268-2806. www.cil.nu-gro.com.

C-I-L Oval® ICI Canada Inc./Lic. Nu-Gro. LARVEX® Zeneca Corp.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))