

95.02.23

RIVERSIDE(R) MCPA ESTER 500

EMULSIFIABLE LIQUID HERBICIDE

FOR THE CONTROL OF BROADLEAF WEEDS
IN BARLEY, WHEAT, RYE AND FLAX.

AGRICULTURAL

READ THE LABEL BEFORE USING

WARNING POISON

GUARANTEE: MCPA(present as mixed butyl ester) 500 g/L

REGISTRATION NO. 23491 PEST CONTROL PRODUCTS ACT

NET CONTENTS: 10 litres

Made in the USA
EPA Est. No. 42750-MO-1

RIVERSIDE/TERRA CORP.
600 FOURTH STREET
SIOUX CITY, Iowa 51101
PHONE: (712) 277-1340

PEEL BACK BOOK HERE AND RESEAL AFTER OPENING

RIVERSIDE SERVES AGRICULTURE. AGRICULTURE SERVES EVERYONE.

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN. DO NOT SPRAY during windy conditions or when air temperature is above 27'C as vapor or spray mist may drift on desirable broadleaf plants. Susceptible plants include beans, sugarbeets, tomatoes, rape, sunflowers, peas, clovers, trees and shrubs. Hazard of drift can be reduced by coarse sprays, low spraying pressure. DO NOT CONTAMINATE domestic or irrigation waters. Thoroughly clean sprayer after

use. DO NOT STORE near or in the same room as seed, fertilizers, insecticides or fungicides. Allow 7 days between application of product and grazing/cutting for hay.

FIRST AID

Avoid contact with skin, eyes and clothing. If swallowed, do not induce vomiting, but rush patient to nearest hospital taking this container or contact a poison control centre. This product contains petroleum distillates. In case of contact on skin, wash with plenty of soap and water; if in eyes, flush with plenty of water and get medical attention.

STORAGE

May be stored at any temperature. Shake well after storing for one year or longer.

DISPOSAL

1. Rinse the emptied container thoroughly and add the rinsings to the spray mixture in the tank.
2. Follow provincial instructions for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.
4. Dispose of the container in accordance with provincial requirements.
5. For information on the disposal of unused, unwanted product and the cleanup of spills, contact the regional office of Environmental Protection, Environment Canada.

NOTICE TO USER

This control product is to be used only in accordance with the directions on this label. It is an offense under the Pest Control Products Act to use a control product under unsafe conditions.

DIRECTIONS FOR USE

This product may be applied by aerial or conventional ground spraying equipment. Aerial spraying is a specialized job, hence

only competent, reliable operators should be employed.

Apply lower rates to weed seedlings in 2 to 4 leaf stage which are growing rapidly and higher rates for more advanced weeds or during dry and cool weather or heavy infestation.

RATE

CROP	L PER Ha	HECTARES TREATED BY 20 LITRES
Barley, Oats, Rye, Wheat	1.25-2.25 L	9-16

DIRECTIONS

Apply sufficient water to obtain good coverage. Treat spring seeded cereals when crop is in 2-4 leaf stage, combining maximum crop tolerance and weed susceptibility. May be applied at early shot blade stage and again from the milk stage to full maturity. For fall seeded wheat or rye, treat when crop growth begins until early shot blade stage. Use lower rates for cereals under-seeded to legumes.

RATE

CROP	L per Ha	HECTARES TREATED BY 20 LITRES
Flax	1.25-2.25 L	9-16

DIRECTIONS

Spray crop when plants are 5 cm high until pre-bud stage.

RATE

CROP	mL per Ha	HECTARES TREATED BY 20 LITRES
Established Alfalfa, Alsike, Red Clover, and Grasses.	700 mL-2.0 L	10-29

DIRECTIONS

Apply 150 to 200 litres of water per hectare. Treat before the development of legumes has occurred and when they are covered by a canopy of weeds and crop. Do not apply in the presence of sweet clover.

RATE

CROP	L per Ha	HECTARES TREATED BY 20 L
UNCROPPED LAND (drainage ditches, pastures, roadsides)	1.75-3.25 L 2.2-4.5 L	6-12 4.5-9

DIRECTIONS

For general weed control, apply 100-200 litres of water per hectare. Treat in early spring or late fall when weeds are in bud. The low rates may be used on more susceptible weeds.

For control of woody plants (poplars, western snowberry and willows). Treat when plants are in full leaf.

NOTICE TO BUYER

Seller's guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons and property arising from the use or handling of this product and accepts the product on that condition.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))