

95.11.03

BOTANIX
LIQUID
LIME SULPHUR
INSECTICIDE/FUNGICIDE
FOR FRUITS, ROSES AND OTHER
ORNAMENTAL PLANTS
FOR BOTH DORMANT AND SUMMER SEASON USE

READ THE LABEL BEFORE USING

DANGER POISON

REGISTRATION NO. 23872
PEST CONTROL PRODUCTS ACT

GUARANTEE:
Sulphide Sulphur 23%

DOMESTIC

1 L

Les Industries Nobel
Boucherville, Québec
J4B 5K1

DIRECTIONS FOR USE:

SHAKE WELL BEFORE USING. Lime Sulphur should mainly be used as a DORMANT SPRAY to clean up overwintering insects and disease infections. During the summer further applications with Lime Sulphur or Botanix Insecticides or Fungicides are necessary to keep plants free from pests. Lime Sulphur may only be mixed with Botanix Dormant Oil, not with any other products. Do not apply during or shortly before temperature reaches 27°C or higher. Shake spray solution while spraying.

Lime Sulphur stains painted wood, brick and stone. Prevent from freezing.

FRUIT TREES: As Dormant Spray, apply before buds break, to

control Blister and Rust Mites, Scale insects, Pear Scale, Red Mite eggs, Aphid eggs, Twig Borer, Plum Back Knot and Peach Leaf Curl. Use 100 mL per 1 litre water.

APPLE AND PEAR: During summer, use 15 mL (1 tablespoon) per litre of water just before and again after blossoming to control powdery mildew, apple and pear scab; repeat 10 to 20 days after end of blossoming. Do not apply on plum in leaf and never on apricot. The apple variety "Delicious" may be injured by summer spraying with Lime Sulphur.

RASPBERRY, BLACKBERRY AND LOGANBERRY: To control rust, anthracnose, cane and spur blight, apply 100 mL in 1 litre of water when buds break.

CURRENT, GOOSEBERRY: To control mildew, rust and scale insects apply 100 mL in 1 litre of water just before buds break. Apply 60 mL (4 tablespoonfuls) in 3 litres of water just before blossoms open and repeat weekly after bloom until fruit colours. May be used up to 1 day before harvest.

ROSES: To control Rose Scale, canker and black spot make dormant application before buds break, using 100 mL in 1 litre of water. For black spot and mildew during summer use 5 mL (1 teaspoonful) in 3 litres of water every one to two weeks.

TREES AND SHRUBS: As Dormant Spray to control scale insects on bark and needles, also Maple gall mite, use 100 mL in 1 litre of water; for Spruce gall aphid use 30 mL (2 tablespoonfuls) in 1 litre of water.

COMBINATION DORMANT SPRAY: Fruit trees, roses and deciduous ornamentals. For overwintering insects, eggs, oystershell scale, San José scale, rose scale, black scale as well as overwintering fungus disease: Mix 120 mL (8 tablespoonfuls) Botanix Lime Sulphur and 60 mL (4 tablespoonfuls) Botanix Dormant Oil in 3 litres of water.

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN. Harmful if swallowed. Avoid contact with eyes and skin. Wash after use. Avoid staining clothing and skin.

FIRST AID: If contact with eyes, rinse with plenty of water for 15 minutes and get medical attention. If contact with skin, wash with water and apply vinegar. If swallowed, give two glasses of water and induce vomiting by inserting finger in throat. Get medical attention immediately or call a poison control centre.

DISPOSAL: Dispose of empty container with household garbage. Do not reuse empty container.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))